

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/82, 15/60, A01H 5/00	A1	(11) International Publication Number: WO 98/58069 (43) International Publication Date: 23 December 1998 (23.12.98)
(21) International Application Number: PCT/US98/12447 (22) International Filing Date: 16 June 1998 (16.06.98) (30) Priority Data: 60/049,955 17 June 1997 (17.06.97) US (71) Applicant: MONSANTO COMPANY [US/US]; 800 N. Lindbergh Boulevard, St. Louis, MO 63167 (US). (72) Inventors: BARRY, Gerard, F.; 5555 Pershing Avenue 2E, St. Louis, MO 63112 (US). CHEIKH, Nordine; 721 Foxwick Drive, Manchester, MO 63011 (US). KISHORE, Ganesh, M.; 15354 Grantley Drive, Chesterfield, MO 63017 (US). (74) Agent: KAMMERER, Patricia, A.; Arnold, White & Durkee, P.O. Box 4433, Houston, TX 77210 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: EXPRESSION OF FRUCTOSE 1,6 BISPHOSPHATE ALDOLASE IN TRANSGENIC PLANTS		
(57) Abstract <p>Fructose-1,6-bisphosphate aldolase (FDA) is an enzyme reversibly catalyzing the reaction converting triosephosphate into fructose-1,6-bisphosphate. In the leaf, this enzyme is located in the chloroplast (starch synthesis) and the cytosol (sucrose biosynthesis). Transgenic plants were generated that express the <i>E. coli fda</i> gene in the chloroplast to improve plant yield by increasing leaf starch biosynthetic ability in particular and sucrose production in general. Leaves from plants expressing the <i>fda</i> transgene showed a significantly higher starch accumulation, as compared to control plants expressing the null vector, particularly early in the photoperiod, but had lower leaf sucrose. Transgenic plants also had a significantly higher root mass. Furthermore, transgenic potatoes expressing <i>fda</i> exhibited improved uniformity of solids.</p>		

BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

EXPRESSION OF FRUCTOSE 1,6 BISPAPHATE

ALDOLASE IN TRANSGENIC PLANTS

This invention relates to the expression of fructose 1,6 bisphosphate aldolase (FDA) in transgenic plants to increase or improve plant growth and development, yield, vigor, stress tolerance, carbon allocation and storage into various storage pools, and distribution of starch. Transgenic plants expressing FDA have increased carbon assimilation, export and storage in plant source and sink organs, which results in growth, yield and quality improvements in crop plants.

Recent advances in genetic engineering have provided the prerequisite tools to transform plants to contain alien (often referred to as "heterologous") or improved endogenous genes. These genes can lead either to an improvement of an already existing pathway in plant tissues or to an introduction of a novel pathway to modify product levels, increase metabolic efficiency, and or save on energy cost to the cell. It is presently possible to produce plants with unique physiological and biochemical traits and characteristics of high agronomic and crop processing importance. Traits that play an essential role in plant growth and development, crop yield potential and stability, and crop quality and composition include enhanced carbon assimilation, efficient carbon storage, and increased carbon export and partitioning.

Atmospheric carbon fixation (photosynthesis) by plants represents the major source of energy to support processes in all living organisms. The primary sites of photosynthetic activity, generally referred to as "source organs", are mature leaves and, to a lesser extent, green stems. The major carbon products of source leaves are starch, which represents the transitory storage form of carbohydrate in the chloroplast, and sucrose, which represents the predominant form of carbon transport in higher plants. Other plant parts named "sink organs" (e.g., roots, fruit, flowers, seeds, tubers, and bulbs) are generally not autotrophic and depend on import of sucrose or other major translocatable carbohydrates for their growth and development. The storage sinks deposit the imported metabolites as sucrose and other oligosaccharides, starch and other polysaccharides, proteins, and triglycerides.

In leaves, the primary products of the Calvin Cycle (the biochemical pathway leading to carbon assimilation) are glyceraldehyde 3-phosphate (G3P) and dihydroxyacetone phosphate (DHAP), also known as triose phosphates (triose-P). The

condensation of G3P and DHAP into fructose 1,6 bisphosphate (FBP) is catalyzed reversibly by the enzyme fructose 1,6 bisphosphate aldolase (FDA), and various isozymes are known. The acidic isoenzyme appears to be chloroplastic and comprises about 85% of the total leaf aldolase activity. The basic isoenzyme is cytosolic. Both isoenzymes appear to be encoded by the nuclear genome and are encoded by different genes (Lebherz et al., 1984).

In the leaf, the chloroplast FDA is an essential enzyme in the Calvin Cycle, where its activity generates metabolites for starch biosynthesis. Removal of more than 40% of the plastidic aldolase enzymatic activity by antisense technology reduced leaf starch accumulation as well as soluble proteins and chlorophyll levels but also reduced plant growth and root formation (Sonnewald et al., 1994). In contrast, the cytosolic FDA is part of the sucrose biosynthetic pathway where it catalyzes the reaction of FBP production. Moreover, cytosolic FDA is also a key enzyme in the glycolytic and gluconeogenesis pathways in both source and sink plant tissues.

In the potato industry, production of higher starch and uniform solids tubers is highly desirable and valuable. The current potato varieties that are used for french fry production, such as Russet Burbank and Shepody, suffer from a non-uniform deposition of solids between the tuber pith (inner core) and the cortex (outer core). French fry strips that are taken from pith tissue are higher in water content when compared to outer cortex french fry strips; cortex tissue typically displays a solids level of twenty-four percent whereas pith tissue typically displays a solids level of seventeen percent. Consequently, in the french fry production process, the pith strips need to be blanched, dried, and par-fried for longer times to eliminate the excess water. Adequate processing of the pith fries results in the over-cooking of fries from the high solids cortex. The blanching, drying, and par frying times of the french fry processor need to be adjusted accordingly to accommodate the low solids pith strips and the high solids cortex strips. A higher solids potato with a more uniform distribution of starch from pith to cortex would allow for a more uniform finished fry product, with higher plant throughput and cost savings due to reduced blanch, dry and par-fry times.

Although various fructose 1,6 bisphosphate aldolases have been previously characterized, it has been discovered that overexpression of the enzyme in a transgenic plant provides advantageous results in the plant such as increasing the assimilation, export

and storage of carbon, increasing the production of oils and/or proteins in the plant and improving tuber solids uniformity.

The present invention provides structural DNA constructs that encode a fructose 1,6 bisphosphate aldolase (FDA) enzyme and that are useful in increasing carbon
5 assimilation, export, and storage in plants.

In accomplishing the foregoing, there is provided, in accordance with one aspect of the present invention, a method of producing genetically transformed plants that have elevated carbon assimilation, storage, export, and improved solids uniformity comprising the steps of:

- 10 (a) Inserting into the genome of a plant a recombinant, double-stranded DNA molecule comprising
 - (i) a promoter that functions in the cells of a target plant tissue,
 - (ii) a structural DNA sequence that causes the production of an RNA sequence that encodes a fructose 1,6 bisphosphate aldolase enzyme,
 - 15 (iii) a 3' non-translated DNA sequence that functions in plant cells to cause transcriptional termination and the addition of polyadenylated nucleotides to the 3' end of the RNA sequence;
- (b) obtaining transformed plant cells; and
- (c) regenerating from transformed plant cells genetically transformed plants that
20 have elevated FDA activity.

In another aspect of the present invention there is provided a recombinant, double-stranded DNA molecule comprising in sequence

- (i) a promoter that functions in the cells of a target plant tissue,
- (ii) a structural DNA sequence that causes the production of an RNA sequence
25 that encodes a fructose 1,6 bisphosphate aldolase enzyme,
- (iii) a 3' non-translated DNA sequence that functions in plant cells to cause transcriptional termination and the addition of polyadenylated nucleotides to the 3' end of the RNA sequence.

In a further aspect of the present invention, the structural DNA sequence that
30 causes the production of an RNA sequence that encodes a fructose 1,6 bisphosphate aldolase enzyme is coupled with a chloroplast transit peptide to facilitate transport of the enzyme to the plastid.

In accordance with the present invention, an improved means for increasing carbon assimilation, storage and export in the source tissues of various plants is provided. Further means of improved carbon accumulation in sinks (such as roots, tubers, seeds, stems, and bulbs) are provided, thus increasing the size of various sinks (larger roots, tubers, etc.) and subsequently increasing yield and crop productivity. The increased carbon availability to these sinks would also improve composition and use efficiency in the sink (oil, protein, starch and/or sucrose production, and/or solids uniformity).

Various advantages may be achieved by the aims of the present invention, including:

First, increasing the expression of the FDA enzyme in the chloroplast would increase the flow of carbon through the Calvin Cycle and increase atmospheric carbon assimilation during early photoperiod. This would result in an increase in photosynthetic efficiency and an increase in chloroplast starch production (a leaf carbon storage form degraded during periods when photosynthesis is low or absent). Both of these responses would lead to an increase in sucrose production by the leaf and a net increase in carbon export during a given photoperiod. This increase in source capacity is a desirable trait in crop plants and would lead to increased plant growth, storage ability, yield, vigor, and stress tolerance.

Second, increasing FDA expression in the cytosol of photosynthetic cells would lead to an increase in sucrose production and export out of source leaves. This increase in source capacity is a desirable trait in crop plants and would lead to increased plant growth, storage ability, yield, vigor, and stress tolerance.

Third, expression of FDA in sink tissues can show several desirable traits, such as increased amino acid and/or fatty acid pools via increases in carbon flux through glycolysis (and thus pyruvate levels) in seeds or other sinks and increased starch levels as result of increased production of glucose 6-phosphate in seeds, roots, stems, and tubers where starch is a major storage nonstructural carbohydrate (reverse glycolysis). This increase in sink strength is a desirable trait in crop plants and would lead to increased plant growth, storage ability, yield, vigor, and stress tolerance.

Fourth, the invention is particularly desirable for use in the commercial production of foods derived from potatoes. Potatoes used for the production of french fries and other products suffer from a non-uniform distribution of solids between the tuber pith (inner

core) and the cortex (outer core). Thus, french fry strips from the pith regions of such tubers have a low solids content and a high water content in comparison to cortex strips from the same tubers. Therefore, the french fry processor attempts to adjust the processing parameters so that the final inner strips are sufficiently cooked while the outer cortex strips are not overcooked. The results of such adjustments, however, are highly variable and may lead to poor quality product. Transgenic potatoes expressing *fda* will provide to the french-fry and potato chip processor a raw product that consistently displays a higher tuber solids uniformity with acceptable agronomic traits. In the french fry plant production process, inner pith fry strips from higher solids uniformity tubers will require less time to blanch, less time to dry to a specific solids content, and less time to par-fry before freezing and shipping to retail and institutional end-users.

Therefore, with respect to potatoes, the present invention provides 1) a higher quality, more uniform finish fry product in which french fries from all tuber regions, when processed, are nearly the same, 2) a higher through-put in the french fry processing plant due to lower processing times, and 3) processor cost savings due to lower energy input required for lower blanch, dry, and par-fry times. A raw tuber product that displays a higher solids uniformity will also produce a potato chip that has a reduced saddle curl, and a reduced tendency for center bubble, which are undesirable qualities in the potato chip industry. Reduced fat content would also result; this would contribute to improved consumer appeal and lower oil use (and costs) for the processor. The increase in solids uniformity will also translate to an increase in overall tuber solids. For both the french fry and chipping industries, this overall tuber solids increase will also result in higher through-put in the processing plant due to lower processing times, and cost savings due to lower energy input for blanching, drying, par-frying, and finish frying.

Figure 1 shows the nucleotide sequence and deduced amino acid sequence of a fructose 1,6 bisphosphate aldolase gene from *E. coli* (SEQ ID No:1).

Figure 2 shows a plasmid map for plant transformation vector pMON17524.

Figure 3 shows a plasmid map for plant transformation vector pMON17542.

Figure 4 shows the change in diurnal fluctuations of sucrose, glucose, and starch levels in tobacco leaves expressing the *fda* transgene (pMON17524) and control (pMON17227). The light period is from 7:00 to 19:00 hours. Only fully expanded and non-senescing leaves were sampled.

5

Figure 5 shows a plasmid map for plant transformation vector pMON13925.

Figure 6 shows a plasmid map for plant transformation vector pMON17590.

10

Figure 7 shows a plasmid map for plant transformation vector pMON13936.

Figure 8 shows a plasmid map for plant transformation vector pMON17581.

15

Figure 9 shows potato tuber cross-sections of improved solids uniformity Segal Russet Burbank lines (top row) versus unimproved nontransgenic Russet Burbank (bottom row).

This invention is directed to a method for producing plant cells and plants demonstrating an increased or improved growth and development, yield, quality, starch storage uniformity, vigor, and/or stress tolerance. The method utilizes a DNA sequence encoding an *fda* (fructose 1,6 bisphosphate aldolase) gene integrated in the cellular genome of a plant as the result of genetic engineering and causes expression of the FDA enzyme in the transgenic plant so produced. Plants that overexpress the FDA enzyme exhibit increased carbon flow through the Calvin Cycle and increased atmospheric carbon assimilation during early photoperiod resulting in an increase in photosynthetic efficiency and an increase in starch production. Thus, such plants exhibit higher levels of sucrose production by the leaf and the ability to achieve a net increase in carbon export during a given photoperiod. This increase in source capacity leads to increased plant growth that in turn generates greater biomass and/or increases the size of the sink and ultimately providing greater yields of the transgenic plant. This greater biomass or increased sink size may be evidenced in different ways or plant parts depending on the particular plant species or growing conditions of the plant overexpressing the FDA enzyme. Thus, increased size resulting from overexpression of FDA may be seen in the seed, fruit, stem,

leaf, tuber, bulb or other plant part depending upon the plant species and its dominant sink during a particular growth phase and upon the environmental effects caused by certain growing conditions, e.g. drought, temperature or other stresses. Transgenic plants overexpressing FDA may therefore have increased carbon assimilation, export and storage
5 in plant source and sink organs, which results in growth, yield, and uniformity and quality improvements.

Plants overexpressing FDA may also exhibit desirable quality traits such as increased production of starch, oils and/or proteins depending upon the plant species overexpressing the FDA. Thus, overexpression of FDA in a particular plant species may
10 affect or alter the direction of the carbon flux thereby directing metabolite utilization and storage either to starch production, protein production or oil production via the role of FDA in the glycolysis and gluconeogenesis metabolic pathways.

The mechanism whereby the expression of exogenous FDA modifies carbon relationships is believed to derive from source-sink relationships. The leaf tissue is a
15 sucrose source, and if more sucrose resulting from the activity of increased FDA expression is transported to a sink, it results in increased storage carbon (sugars, starch, oil, protein, etc.) or nitrogen (protein, etc.) per given weight of the sink tissue.

The expression in a plant of a gene that exists in double-stranded DNA form involves transcription of messenger RNA (mRNA) from one strand of the DNA by RNA
20 polymerase enzyme, and the subsequent processing of the mRNA primary transcript inside the nucleus. This processing involves a 3' non-translated region, which adds polyadenylate nucleotides to the 3' end of the RNA. Transcription of DNA into mRNA is regulated by a region of DNA usually referred to as the promoter. The promoter region contains a sequence of bases that signals RNA polymerase to associate with the DNA and
25 to initiate the transcription of mRNA using one of the DNA strands as a template to make a corresponding complimentary strand of RNA. This RNA is then used as a template for the production of the protein encoded therein by the cells protein biosynthetic machinery.

A number of promoters that are active in plant cells have been described in the literature. These include the nopaline synthase (NOS) and octopine synthase (OCS)
30 promoters (which are carried on tumor-inducing plasmids of *Agrobacterium tumefaciens*), the caulimovirus promoters such as the cauliflower mosaic virus (CaMV) 19S and 35S and the figwort mosaic virus (FMV) 35S-promoters, the light-inducible promoter from the

small subunit of ribulose-1,5-bisphosphate carboxylase (ssRUBISCO), a very abundant plant polypeptide, and the chlorophyll a/b binding protein gene promoters, etc. All of these promoters have been used to create various types of DNA constructs that have been expressed in plants; see, e.g., PCT publication WO 84/02913.

5 Promoters that are known to or are found to cause transcription of DNA in plant cells can be used in the present invention. Such promoters may be obtained from a variety of sources such as plants and plant viruses and include, but are not limited to, the enhanced CaMV35S promoter and promoters isolated from plant genes such as ssRUBISCO genes. As described below, it is preferred that the particular promoter selected should be capable
10 of causing sufficient expression to result in the production of an effective amount of fructose 1,6 bisphosphate aldolase enzyme to cause the desired increase in carbon assimilation, export or storage. Expression of the double-stranded DNA molecules of the present invention can be driven by a constitutive promoter, expressing the DNA molecule in all or most of the tissues of the plant. Alternatively, it may be preferred to cause
15 expression of the *fda* gene in specific tissues of the plant, such as leaf, stem, root, tuber, seed, fruit, etc. The promoter chosen will have the desired tissue and developmental specificity. Those skilled in the art will recognize that the amount of fructose 1,6 bisphosphate aldolase needed to induce the desired increase in carbon assimilation, export, or storage may vary with the type of plant. Therefore, promoter function should be
20 optimized by selecting a promoter with the desired tissue expression capabilities and approximate promoter strength and selecting a transformant that produces the desired fructose 1,6 bisphosphate aldolase activity or the desired change in metabolism of carbohydrates in the target tissues. This selection approach from the pool of transformants is routinely employed in expression of heterologous structural genes in plants because
25 there is variation between transformants containing the same heterologous gene due to the site of gene insertion within the plant genome (commonly referred to as "position effect"). In addition to promoters that are known to cause transcription (constitutively or tissue-specific) of DNA in plant cells, other promoters may be identified for use in the current invention by screening a plant cDNA library for genes that are selectively or preferably
30 expressed in the target tissues of interest and then isolating the promoter regions by methods known in the art. In particular, it may be desirable to use a bundle sheath cell specific (or cell enhanced expression) promoter for use with C4 plants such as corn,

sorghum, and sugarcane to obtain the yield benefits of overexpression of FDA and not use a constitutive promoter or a promoter with mesophyll cell enhanced expression properties.

For the purpose of expressing the *fda* gene in source tissues of the plant, such as the leaf or stem, it is preferred that the promoters utilized in the double-stranded DNA molecules of the present invention have relatively high expression in these specific tissues. For this purpose, one may also choose from a number of promoters for genes with leaf-specific or leaf-enhanced expression. Examples of such genes known from the literature are the chloroplast glutamine synthetase GS2 from pea (Edwards et al., 1990), the chloroplast fructose-1,6-bisphosphatase (FBPase) from wheat (Lloyd et al., 1991), the nuclear photosynthetic ST-LS1 from potato (Stockhaus et al., 1989), and the phenylalanine ammonia-lyase (PAL) and chalcone synthase (CHS) genes from *Arabidopsis thaliana* (Leyva et al., 1995). Also shown to be active in photosynthetically active tissues are the ribulose-1,5-bisphosphate carboxylase (RUBISCO), isolated from eastern larch (*Larix laricina*) (Campbell et al., 1994); the *cab* gene, encoding the chlorophyll a/b-binding protein of PSII, isolated from pine (*cab6*; Yamamoto et al., 1994), wheat (*Cab-1*; Fejes et al., 1990), spinach (*CAB-1*; Luebbertstedt et al., 1994), and rice (*cab1R*; Luan et al., 1992); the pyruvate orthophosphate dikinase (PPDK) from maize (Matsuoka et al., 1993); the tobacco Lhcb1*2 gene (Cerdan et al., 1997); the *Arabidopsis thaliana* SUC2 sucrose-H⁺ symporter gene (Truernit et al., 1995); and the thylacoid membrane proteins, isolated from spinach (*psaD*, *psaF*, *psaE*, *PC*, *FNR*, *atpC*, *atpD*, *cab*, *rbcS*; Oelmueller et al., 1992). Other chlorophyll a/b-binding proteins have been studied and described in the literature, such as LhcB and PsbP from white mustard (*Sinapis alba*; Kretsch et al., 1995). Homologous promoters to those described here may also be isolated from and tested in the target or related crop plant by standard molecular biology procedures.

For the purpose of expressing the *fda* in sink tissues of the plant, for example the tuber of the potato plant; the fruit of tomato; or seed of maize, wheat, rice, or barley, it is preferred that the promoters utilized in the double-stranded DNA molecules of the present invention have relatively high expression in these specific tissues. A number of genes with tuber-specific or tuber-enhanced expression are known, including the class I patatin promoter (Bevan et al., 1986; Jefferson et al., 1990); the potato tuber ADPGPP genes, both the large and small subunits (Muller et al., 1990); sucrose synthase (Salañoubat and Belliard, 1987, 1989); the major tuber proteins including the 22 kDa protein complexes

and proteinase inhibitors (Hannapel, 1990); the granule bound starch synthase gene (GBSS) (Rohde et al., 1990); and the other class I and II patatins (Rocha-Sosa et al., 1989; Mignery et al., 1988). Other promoters can also be used to express a fructose 1,6 bisphosphate aldolase gene in specific tissues, such as seeds or fruits. The promoter for β -conglycinin (Tierney, 1987) or other seed-specific promoters, such as the napin and phaseolin promoters, can be used to over-express an *fda* gene specifically in seeds. The zeins are a group of storage proteins found in maize endosperm. Genomic clones for zein genes have been isolated (Pedersen et al., 1982), and the promoters from these clones, including the 15 kDa, 16 kDa, 19 kDa, 22 kDa, 27 kDa, and gamma genes, could also be used to express an *fda* gene in the seeds of maize and other plants. Other promoters known to function in maize, wheat, or rice include the promoters for the following genes: *waxy*, *Brittle*, *Shrunken 2*, branching enzymes I and II, starch synthases, debranching enzymes, oleosins, glutelins, and sucrose synthases. Particularly preferred promoters for maize endosperm expression, as well as in wheat and rice, of an *fda* gene is the promoter for a glutelin gene from rice, more particularly the Osgt-1 promoter (Zheng et al., 1993); the maize granule-bound starch synthase (*waxy*) gene (zmGBS); the rice small subunit ADPGPP promoter (osAGP); and the zein promoters, particularly the maize 27 kDa zein gene promoter (zm27) (see, generally, Russell et al., 1997). Examples of promoters suitable for expression of an *fda* gene in wheat include those for the genes for the ADPGlucose pyrophosphorylase (ADPGPP) subunits, for the granule bound and other starch synthases, for the branching and debranching enzymes, for the embryogenesis-abundant proteins, for the gliadins, and for the glutenins. Examples of such promoters in rice include those for the genes for the ADPGPP subunits, for the granule bound and other starch synthases, for the branching enzymes, for the debranching enzymes, for sucrose synthases, and for the glutelins. A particularly preferred promoter is the promoter for rice glutelin, Osgt-1. Examples of such promoters for barley include those for the genes for the ADPGPP subunits, for the granule bound and other starch synthases, for the branching enzymes, for the debranching enzymes, for sucrose synthases, for the hordeins, for the embryo globulins, and for the aleurone-specific proteins.

The solids content of root tissue may be increased by expressing an *fda* gene behind a root-specific promoter. An example of such a promoter is the promoter from the acid chitinase gene (Samac et al., 1990). Expression in root tissue could also be

accomplished by utilizing the root-specific subdomains of the CaMV35S promoter that have been identified (Benfey et al., 1989).

The RNA produced by a DNA construct of the present invention may also contain a 5' non-translated leader sequence. This sequence can be derived from the promoter selected to express the gene and can be specifically modified so as to increase translation of the mRNA. The 5' non-translated regions can also be obtained from viral RNAs, from suitable eukaryotic genes, or from a synthetic gene sequence. The present invention is not limited to constructs, as presented in the following examples, wherein the non-translated region is derived from the 5' non-translated sequence that accompanies the promoter sequence. Rather, the non-translated leader sequence can be derived from an unrelated promoter or coding sequence.

In monocots, an intron is preferably included in the gene construct to facilitate or enhance expression of the coding sequence. Examples of suitable introns include the HSP70 intron and the rice actin intron, both of which are known in the art. Another suitable intron is the castor bean catalase intron (Suzuki et al., 1994)

Polyadenylation signal

The 3' non-translated region of the chimeric plant gene contains a polyadenylation signal that functions in plants to cause the addition of polyadenylate nucleotides to the 3' end of the RNA. Examples of suitable 3' regions are (1) the 3' transcribed, non-translated regions containing the polyadenylation signal of *Agrobacterium* tumor-inducing (Ti) plasmid genes, such as the nopaline synthase (NOS) gene, and (2) plant genes like the soybean storage protein genes and the small subunit of the ribulose-1,5-bisphosphate carboxylase (ssRUBISCO) gene.

Plastid-directed expression of fructose-1,6-bisphosphate aldolase activity

In one embodiment of the invention, the *fda* gene may be fused to a chloroplast transit peptide, in order to target the FDA protein to the plastid. As used hereinafter, chloroplast and plastid are intended to include the various forms of plastids including amyloplasts. Many plastid-localized proteins are expressed from nuclear genes as precursors and are targeted to the plastid by a chloroplast transit peptide (CTP), which is removed during the import steps. Examples of such chloroplast proteins include the small subunit of ribulose-1,5-bisphosphate carboxylase (ssRUBISCO, SSU), 5-enolpyruvateshikimate-3-phosphate synthase (EPSPS), ferredoxin, ferredoxin

oxidoreductase, the light-harvesting-complex protein I and protein II, and thioredoxin F. It has been demonstrated that non-plastid proteins may be targeted to the chloroplast by use of protein fusions with a CTP and that a CTP sequence is sufficient to target a protein to the plastid. Those skilled in the art will also recognize that various other chimeric
5 constructs can be made that utilize the functionality of a particular plastid transit peptide to import the fructose-1,6-diphosphate aldolase enzyme into the plant cell plastid. The *fda* gene could also be targeted to the plastid by transformation of the gene into the chloroplast genome (Daniell et al., 1998).

Fructose 1,6 biphosphate aldolases

10 As used herein, the term "fructose 1, 6-bisphosphate aldolase" means an enzyme (E.C. 4.1.2.13) that catalyzes the reversible cleavage of fructose 1,6-bisphosphate to form glyceraldehyde 3-phosphate (G3P) and dihydroxyacetone phosphate (DHAP). Aldolase enzymes are divided into two classes, designated class I and class II (Witke and Gotz, 1993). Various *fda* genes encoding the enzyme have been sequenced, as have numerous
15 proteins, such as the cytosolic enzyme from maize (GenBank Accession S07789;S10638), cytosolic enzyme from rice (GenBank Accession JQ0543), cytosolic enzyme from spinach (GenBank Accession S31091;S22093), from *Arabidopsis thaliana* (GenBank Accession S11958), from spinach chloroplast (GenBank Accession S31090;A21815;S22092), from yeast (*S. cerevisiae*) (GenBank Accession S07855; S37882; S12945; S39178;
20 S44523;X75781), from *Rhodobacter sphaeroides* (GenBank Accession B40767;D41080), from *B. subtilis* (GenBank Accession S55426; D32354; E32354; D41835), from garden pea (GenBank Accession S29048; S34411), from garden pea chloroplast (GenBank Accession S29047; S34410), from maize (GenBank Accession S05019), from *Chlamydomonas reinhardtii* (GenBank Accession S48639; S58485; S58486; S34367),
25 from *Corynebacterium glutamicum* (GenBank Accession S09283; X17313), from *Campylobacter jejuni* (GenBank Accession S52413), from *Haemophilus influenzae* (strain Rd KW20) (GenBank Accession C64074), from *Streptococcus pneumonia* (GenBank Accession AJ005697), from rice (GenBank Accession X53130), and from the maize anaerobically regulated gene (GenBank Accession X12872).

30 The class I enzymes may be isolated from higher eukaryotes, such as animals and plants, and in some prokaryotes, including *Peptococcus aerogens*, (Lebherz and Rutter, 1973), *Lactobacillus casei* (London and Kline, 1973), *Escherichia coli* (Stribling and

Perham, 1973), *Mycobacterium smegmatis* (Bai et al., 1975), and most staphylococcal species (Gotz et al., 1979). The gene for the FDA enzyme may be obtained by known methods and has already been done so for several organisms, such as rabbit (Lai et al., 1974), human (Besmond et al., 1983), rat (Tsutsumi et al., 1984), *Trypanosoma brucei* 5 (Clayton, 1985), and *Arabidopsis thaliana* (Chopra et al., 1990). These class I enzymes are invariably tetrameric proteins with a total molecular weight of about 160 kDa and function by imine formation between the substrate and a lysine residue in the active site (Alfounder et al., 1989).

In animal, three class I isozymes, classified as A, B, and C, are expressed in the 10 cytosol of muscle, liver, and brain tissue respectively, and they differ from plant aldolases in their expression and compartmentation patterns (Joh et al., 1986). In the leaves of higher plants, FDA is a class I enzyme, and two different isoenzymes within the class have been documented. One is contained in the chloroplast and the other in the cytosol (Lebherz et al., 1984). The acidic plant isozyme appear to be chloroplastic and comprises 15 about 85% of the total leaf aldolase activity. The basic plant isozyme is cytosolic, and both isozymes appear to be encoded by the nuclear genome and are encoded by different genes (Lebherz et al., 1984).

The class II type aldolases are normally dimeric with molecular mass of approximately 80 kDa, and their activity depends on divalent metal ions. The class II 20 enzymes may be isolated from prokaryotes, such as blue-green algae and bacteria, and eukaryotic green algae and fungi (Baldwin et al., 1978). The gene for the FDA class II enzyme may be obtained by known methods and has already been done so from several organisms including *Saccharomyces cerevisiae* (Jack and Harris, 1971), *Bacillus stearothermophilus* (Jack, 1973), and *Escherichia coli* (Baldwin et al., 1978).

25 It is believed that highly homologous class II fructose 1, 6-bisphosphate aldolases with similar catalyzing activity will also be found in other species of microorganism, such as *Saccharomyces* (*Saccharomyces cerevisiae*); *Bacillus* (*Bacillus subtilis*); *Rhodobacter* (*Rhodobacter sphaeroides*); *Plasmodium* (*Plasmodium falciparum*, *Plasmodium berghei*); *Trypanosoma* (*Trypanosoma brucei*); *Chlamydomonas* (*Chlamydomonas reinhardtii*); 30 *Candida* (*Candida albicans*); *Corynebacterium* (*Corynebacterium glutamicum*); *Campylobacter* (*Campylobacter jejuni*); and *Haemophilus* (*Haemophilus influenza*).

Such sequences can be readily isolated by methods well known in the art, for example by nucleic acid hybridization. The hybridization properties of a given pair of nucleic acids are an indication of their similarity or identity. Nucleic acid sequences can be selected on the basis of their ability to hybridize with known *fda* sequences. Low stringency conditions may be used to select sequences with less homology or identity. One may wish to employ conditions such as about 0.15 M to about 0.9 M sodium chloride, at temperatures ranging from about 20°C to about 55°C. High stringency conditions may be used to select for nucleic acid sequences with higher degrees of identity to the disclosed sequences. Conditions typically employed may include about 0.02 M to about 0.15 M sodium chloride, about 0.5% to about 5% casein, about 0.02% SDS or about 0.1% N-laurylsarcosine, about 0.001 M to about 0.03 M sodium citrate, at hybridization temperatures between about 50°C and about 70°C. More preferably, high stringency conditions are about 0.02 M sodium chloride, about 0.5% casein, about 0.02% SDS, about 0.001 M sodium citrate, at a temperature of about 50°C. The skilled individual will recognize that numerous variations are possible in the conditions and means by which nucleic acid hybridization can be performed to isolate *fda* sequences having similarity to *fda* sequences known in the art and are not limited to those explicitly disclosed herein. Preferably, such an approach is used to isolate *fda* sequences having greater than about 60% identity with the disclosed *E.coli fda* sequence, more preferably greater than about 70% identity, most preferably greater than about 80% identity.

Depending on growth conditions *Euglena gracilis*, *Chlamydomonas mundana*, and *Chlamydomonas reinhardtii* produce either a class I or a class II aldolase (Cremona, 1968; Russell and Gibbs, 1967; Guerrini et al., 1971).

The isolation of a class II *fda* gene from *E.coli* is described in the following examples. Its DNA sequence is given as SEQ ID NO:1 and shown in Figure 1. The amino acid sequence is shown in SEQ ID NO:2 and shown in Figure 1. This gene can be used as isolated by inserting it into plant expression vectors suitable for the transformation method of choice as described. The *E. coli* FDA enzyme has an apparent pH optimum range near pH 7-9 and retains activity in the lower pH range of 5-7 (Baldwin et al., 1978; Alfounder et al., 1989).

Thus, many different genes that encode a fructose 1,6 bisphosphate aldolase activity may be isolated and used in the present invention.

Synthetic gene construction

A carbohydrate metabolizing enzyme considered in this invention includes any sequence of amino acids, such as protein, polypeptide, or peptide fragment, that demonstrates the ability to catalyze a reaction involved in the synthesis or degradation of starch or sucrose. These can be sequences obtained from a heterologous source, such as algae, bacteria, fungi, and protozoa, or endogenous plant sequences, by which is meant any sequence that can be naturally found in a plant cell, including native (indigenous) plant sequences as well as sequences from plant viruses or plant pathogenic bacteria.

It will be recognized by one of ordinary skill in the art that carbohydrate metabolizing enzyme gene sequences may also be modified using standard techniques such as site-specific mutation or PCR, or modification of the sequence may be accomplished by producing a synthetic nucleic acid sequence and will still be considered a carbohydrate biosynthesis enzyme nucleic acid sequence of this invention. For example, "wobble" positions in codons may be changed such that the nucleic acid sequence encodes the same amino acid sequence, or alternatively, codons can be altered such that conservative amino acid substitutions result. In either case, the peptide or protein maintains the desired enzymatic activity and is thus considered part of this invention.

A nucleic acid sequence to a carbohydrate metabolizing enzyme may be a DNA or RNA sequence, derived from genomic DNA, cDNA, mRNA, or may be synthesized in whole or in part. The structural gene sequences may be cloned, for example, by isolating genomic DNA from an appropriate source and amplifying and cloning the sequence of interest using a polymerase chain reaction (PCR). Alternatively, the gene sequences may be synthesized, either completely or in part, especially where it is desirable to provide plant-preferred sequences. Thus, all or a portion of the desired structural gene may be synthesized using codons preferred by a selected plant host. Plant-preferred codons may be determined, for example, from the codons used most frequently in the proteins expressed in a particular plant host species. Other modifications of the gene sequences may result in mutants having slightly altered activity.

If desired, the gene sequence of the *fda* gene can be changed without changing the protein sequence in such a manner as may increase expression and thus even more positively affect carbohydrate content in transformed plants. A preferred manner for making the changes in the gene sequence is set out in PCT Publication WO 90/10076. A

gene synthesized by following the methodology set out therein may be introduced into plants as described below and result in higher levels of expression of the FDA enzyme. This may be particularly useful in monocots such as maize, rice, wheat, sugarcane, and barley.

5 Combinations with other transgenes

 The effect of *fda* in transgenic plants may be enhanced by combining it with other genes that positively affect carbohydrate assimilation or content, such as a gene encoding for a sucrose phosphorylase as described in PCT Publication WO 96/24679, or ADPGPP genes such as the *E. coli glgC* gene and its mutant *glgC16*. PCT Publication WO
10 91/19806 discloses how to incorporate the latter gene into many plant species in order to increase starch or solids. Another gene that can be combined with *fda* to increase carbon assimilation, export or storage is a gene encoding for sucrose phosphate synthase (SPS). PCT Publication WO 92/16631 discloses one such gene and its use in transgenic plants.
 Plant transformation/regeneration

15 In developing the nucleic acid constructs of this invention, the various components of the construct or fragments thereof will normally be inserted into a convenient cloning vector, e.g., a plasmid that is capable of replication in a bacterial host, e.g., *E. coli*. Numerous vectors exist that have been described in the literature, many of which are commercially available. After each cloning, the cloning vector with the desired insert may
20 be isolated and subjected to further manipulation, such as restriction digestion, insertion of new fragments or nucleotides, ligation, deletion, mutation, resection, etc. so as to tailor the components of the desired sequence. Once the construct has been completed, it may then be transferred to an appropriate vector for further manipulation in accordance with the manner of transformation of the host cell.

25 A recombinant DNA molecule of the invention typically includes a selectable marker so that transformed cells can be easily identified and selected from non-transformed cells. Examples of such include, but are not limited to, a neomycin phosphotransferase (*nptII*) gene (Potrykus et al., 1985), which confers kanamycin resistance. Cells expressing the *nptII* gene can be selected using an appropriate antibiotic
30 such as kanamycin or G418. Other commonly used selectable markers include the *bar* gene, which confers bialaphos resistance; a mutant EPSP synthase gene (Hinchee et al., 1988), which confers glyphosate resistance; a nitrilase gene, which confers resistance to

bromoxynil (Stalker et al., 1988); a mutant acetolactate synthase gene (ALS), which confers imidazolinone or sulphonylurea resistance (European Patent Application 154,204, 1985); and a methotrexate resistant DHFR gene (Thillet et al., 1988).

Plants that can be made to have enhanced carbon assimilation, increased carbon export and partitioning by practice of the present invention include, but are not limited to, *Acacia*, alfalfa, aneth, apple, apricot, artichoke, arugula, asparagus, avocado, banana, barley, beans, beet, blackberry, blueberry, broccoli, brussels sprouts, cabbage, canola, cantaloupe, carrot, cassava, cauliflower, celery, cherry, cilantro, citrus, clementines, coffee, corn, cotton, cucumber, Douglas fir, eggplant, endive, escarole, eucalyptus, fennel, 5
10 figs, gourd, grape, grapefruit, honey dew, jicama, kiwifruit, lettuce, leeks, lemon, lime, Loblolly pine, mango, melon, mushroom, nut, oat, oil seed rape, okra, onion, orange, an ornamental plant, papaya, parsley, pea, peach, peanut, pear, pepper, persimmon, pine, pineapple, plantain, plum, pomegranate, poplar, potato, pumpkin, quince, radiata pine, radicchio, radish, raspberry, rice, rye, sorghum, Southern pine, soybean, spinach, squash, 15
strawberry, sugarbeet, sugarcane, sunflower, sweet potato, sweetgum, tangerine, tea, tobacco, tomato, triticale, turf, a vine, watermelon, wheat, yams, and zucchini.

A double-stranded DNA molecule of the present invention containing an *fda* gene can be inserted into the genome of a plant by any suitable method. Suitable plant transformation vectors include those derived from a Ti plasmid of *Agrobacterium* 20
tumefaciens, as well as those disclosed, e.g., by Herrera-Estrella et al. (1983), Bevan (1984), Klee et al. (1985) and EPO publication 120,516. In addition to plant transformation vectors derived from the Ti or root-inducing (Ri) plasmids of *Agrobacterium*, alternative methods can be used to insert the DNA constructs of this invention into plant cells. Such methods may involve, for example, the use of liposomes, 25
electroporation, chemicals that increase free DNA uptake, free DNA delivery via microprojectile bombardment, and transformation using viruses or pollen. DNA may also be inserted into the chloroplast genome (Daniell et al., 1998).

A plasmid expression vector suitable for the introduction of an *fda* gene in monocots using microprojectile bombardment is composed of the following: a promoter 30
that is specific or enhanced for expression in the starch storage tissues in monocots, generally the endosperm, such as promoters for the zein genes found in the maize endosperm (Pedersen et al., 1982); an intron that provides a splice site to facilitate

expression of the gene, such as the Hsp70 intron (PCT Publication WO93/19189); and a 3' polyadenylation sequence such as the nopaline synthase 3' sequence (NOS 3'; Fraley et al., 1983). This expression cassette may be assembled on high copy replicons suitable for the production of large quantities of DNA.

5 A particularly useful *Agrobacterium*-based plant transformation vector for use in transformation of dicotyledonous plants is plasmid vector pMON530 (Rogers et al., 1987). Plasmid pMON530 is a derivative of pMON505 prepared by transferring the 2.3 kb StuI-HindIII fragment of pMON316 (Rogers et al., 1987) into pMON526. Plasmid pMON526 is a simple derivative of pMON505 in which the SmaI site is removed by digestion with
10 XmaI, treatment with Klenow polymerase and ligation. Plasmid pMON530 retains all the properties of pMON505 and the CaMV35S-NOS expression cassette and now contains a unique cleavage site for SmaI between the promoter and polyadenylation signal.

Binary vector pMON505 is a derivative of pMON200 (Rogers et al., 1987) in which the Ti plasmid homology region, LIH, has been replaced with a 3.8 kb HindIII to
15 SmaI segment of the mini RK2 plasmid, pTJS75 (Schmidhauser and Helinski, 1985). This segment contains the RK2 origin of replication, oriV, and the origin of transfer, oriT, for conjugation into *Agrobacterium* using the tri-parental mating procedure (Horsch and Klee, 1986). Plasmid pMON505 retains all the important features of pMON200 including the synthetic multi-linker for insertion of desired DNA fragments, the chimeric
20 NOS/NPTII'/NOS gene for kanamycin resistance in plant cells, the spectinomycin/streptomycin resistance determinant for selection in *E. coli* and *A. tumefaciens*, an intact nopaline synthase gene for facile scoring of transformants and inheritance in progeny, and a pBR322 origin of replication for ease in making large
amounts of the vector in *E. coli*. Plasmid pMON505 contains a single T-DNA border
25 derived from the right end of the pTiT37 nopaline-type T-DNA. Southern blot analyses have shown that plasmid pMON505 and any DNA that it carries are integrated into the plant genome, that is, the entire plasmid is the T-DNA that is inserted into the plant
genome. One end of the integrated DNA is located between the right border sequence and the nopaline synthase gene and the other end is between the border sequence and the
30 pBR322 sequences.

Another particularly useful Ti plasmid cassette vector is pMON17227. This vector is described in PCT Publication WO 92/04449 and contains a gene encoding an enzyme

conferring glyphosate resistance (denominated CP4), which is an excellent selection marker gene for many plants, including potato and tomato. The gene is fused to the *Arabidopsis* EPSPS chloroplast transit peptide (CTP2) and expressed from the FMV promoter as described therein.

5 When adequate numbers of cells (or protoplasts) containing the *fda* gene or cDNA are obtained, the cells (or protoplasts) are regenerated into whole plants. Choice of methodology for the regeneration step is not critical, with suitable protocols being available for hosts from Leguminosae (alfalfa, soybean, clover, etc.), Umbelliferae (carrot, celery, parsnip), Cruciferae (cabbage, radish, canola/rapeseed, etc.), Cucurbitaceae
10 (melons and cucumber), Gramineae (wheat, barley, rice, maize, etc.), Solanaceae (potato, tobacco, tomato, peppers), various floral crops, such as sunflower, and nut-bearing trees, such as almonds, cashews, walnuts, and pecans. See, e.g., Ammirato et al. (1984); Shimamoto et al. (1989); Fromm (1990); Vasil et al. (1990); Vasil et al. (1992); Hayashimoto (1990); and Datta et al. (1990).

15 The following definitions are provided in order to aid those skilled in the art in understanding the detailed description of the present invention.

 The term "promoter" or "promoter region" refers to a nucleic acid sequence, usually found upstream (5') to a coding sequence, that controls expression of the coding sequence by controlling production of messenger RNA (mRNA) by providing the
20 recognition site for RNA polymerase or other factors necessary for start of transcription at the correct site. As contemplated herein, a promoter or promoter region includes variations of promoters derived by means of ligation to various regulatory sequences, random or controlled mutagenesis, and addition or duplication of enhancer sequences. The promoter region disclosed herein, and biologically functional equivalents thereof, are
25 responsible for driving the transcription of coding sequences under their control when introduced into a host as part of a suitable recombinant vector, as demonstrated by its ability to produce mRNA.

 "Regeneration" refers to the process of growing a plant from a plant cell (e.g., plant protoplast or explant).

30 "Transformation" refers to a process of introducing an exogenous nucleic acid sequence (e.g., a vector, recombinant nucleic acid molecule) into a cell or protoplast in

which that exogenous nucleic acid is incorporated into a chromosome or is capable of autonomous replication.

A "transformed cell" is a cell whose DNA has been altered by the introduction of an exogenous nucleic acid molecule into that cell.

5 The term "gene" refers to chromosomal DNA, plasmid DNA, cDNA, synthetic DNA, or other DNA that encodes a peptide, polypeptide, protein, or RNA molecule, and regions flanking the coding sequence involved in the regulation of expression.

"Identity" refers to the degree of similarity between two nucleic acid or protein sequences. An alignment of the two sequences is performed by a suitable computer
10 program. A widely used and accepted computer program for performing sequence alignments is CLUSTALW v1.6 (Thompson et al., 1994). The number of matching bases or amino acids is divided by the total number of bases or amino acids and multiplied by 100 to obtain a percent identity. For example, if two 580 base pair sequences had 145 matched bases, they would be 25 percent identical. If the two compared sequences are of
15 different lengths, the number of matches is divided by the shorter of the two lengths. For example, if there were 100 matched amino acids between 200 and a 400 amino acid proteins, they are 50 percent identical with respect to the shorter sequence. If the shorter sequence is less than 50 bases or amino acids in length, the number of matches are divided by 50 and multiplied by 100 to obtain a percent identity.

20 "C-terminal region" refers to the region of a peptide, polypeptide, or protein chain from the middle thereof to the end that carries the amino acid having a free carboxyl group.

The phrase "DNA segment heterologous to the promoter region" means that the coding DNA segment does not exist in nature in the same gene with the promoter to which
25 it is now attached.

The term "encoding DNA" refers to chromosomal DNA, plasmid DNA, cDNA, or synthetic DNA that encodes any of the enzymes discussed herein.

The term "genome" as it applies to bacteria encompasses both the chromosome and plasmids within a bacterial host cell. Encoding DNAs of the present invention introduced
30 into bacterial host cells can therefore be either chromosomally integrated or plasmid-localized. The term "genome" as it applies to plant cells encompasses not only chromosomal DNA found within the nucleus, but organelle DNA found within subcellular

components of the cell. DNAs of the present invention introduced into plant cells can therefore be either chromosomally integrated or organelle-localized.

The terms "microbe" or "microorganism" refer to algae, bacteria, fungi, and protozoa.

5 The term "mutein" refers to a mutant form of a peptide, polypeptide, or protein.

"N-terminal region" refers to the region of a peptide, polypeptide, or protein chain from the amino acid having a free amino group to the middle of the chain.

"Overexpression" refers to the expression of a polypeptide or protein encoded by a DNA introduced into a host cell, wherein said polypeptide or protein is either not normally
10 present in the host cell, or wherein said polypeptide or protein is present in said host cell at a higher level than that normally expressed from the endogenous gene encoding said polypeptide or protein.

The term "plastid" refers to the class of plant cell organelles that includes amyloplasts, chloroplasts, chromoplasts, elaioplasts, eoplasts, etioplasts, leucoplasts, and
15 proplastids. These organelles are self-replicating and contain what is commonly referred to as the "chloroplast genome," a circular DNA molecule that ranges in size from about 120 kb to about 217 kb, depending upon the plant species, and which usually contains an inverted repeat region.

The phrase "simple carbohydrate substrate" means a monosaccharide or an
20 oligosaccharide but not a polysaccharide; simple carbohydrate substrate includes glucose, fructose, sucrose, lactose. More complex carbohydrate substrates commonly used in media such as corn syrup, starch, and molasses can be broken down to simple carbohydrate substrates.

The term "solids" refers to the nonaqueous component of a tuber (such as in
25 potato) or a fruit (such as in tomato) comprised mostly of starch and other polysaccharides, simple carbohydrates, nonstructural carbohydrates, amino acids, and other organic molecules.

The following examples are provided to better elucidate the practice of the present invention and should not be interpreted in any way to limit the scope of the present
30 invention. Those skilled in the art will recognize that various modifications, truncations, etc., can be made to the methods and genes described herein while not departing from the spirit and scope of the present invention.

EXAMPLES

EXAMPLE 1

cDNA cloning and overexpression

Unless otherwise stated, basic DNA manipulations and genetic techniques, such as PCR, agarose electrophoresis, restriction digests, ligations, *E. coli* transformations, colony screens, and Western blots were performed essentially by the protocols described in Sambrook et al. (1989) or Maniatis et al. (1982).

The *E. coli fda* gene sequence (SEQ ID NO: 1) was obtained from Genbank (Accession Number X14682) and nucleotide primers with homology to the 5' and 3' end were designed for PCR amplification. *E. coli* chromosomal DNA was extracted and the *E. coli fda* gene was amplified by PCR using the 5' oligonucleotide 5'GGGGCCATGGCTAAGATTTTTGATTTTCGTA3' (SEQ ID NO:3) and the 3' oligonucleotide 5'CCCCGAGCTCTTACAGAACGTCGATCGCGTTCAG3' (SEQ ID NO:4). The PCR cycling conditions were as follows: 94°C, 5 min (1 cycle); addition of polymerase; 94°C, 1 min., 60°C, 1 min., 72°C, 2 min.30 sec. (35 cycles). The 1.08 kb PCR product was gel purified and ligated into an *E. coli* expression vector, pMON5723, to form a vector construct that was used for transformation of frozen competent *E. coli* JM101 cells. The pMON5723 vector contains the *E. coli* recA promoter and the T7 gene10 leader (G10L) sequences, which enable high level expression in *E. coli* (Wong et al., 1988). After induction of the transformed cells, a distinct protein band of about 40 kDa was apparent on an SDS PAGE gel, which correlates with the size of the subunit polypeptide chain of the dimeric aldolase II. It was shown that most of the induced protein was present in the soluble phase. A gel slice containing the highly induced protein was isolated and antibodies were produced in a goat, which was injected with the homogenized gel slice (emulsified in Freund's complete adjuvant).

The *fda* gene sequence was subsequently cloned into another *E. coli* expression vector, under the control of the *taq* promoter. Induction with IPTG of JM101 cells transformed with this vector showed the same 40 kDa overexpressed protein band. This new clone was used in an enzyme assay for FDA activity. Cells transformed with this vector construct were grown in a liquid culture, induced with IPTG, and grown for another 3 hours. Subsequently, a 3 mL cell culture was spun down, dissolved in 100mM Tris and sonicated. The cell pellet was spun down, and the crude cell extract supernatant was

assayed for FDA activity, using a coupled enzymatic assay as described by Baldwin et al. (1978). This assay was routinely performed at 30°C.

The reaction was performed in a 1 mL final volume in excess presence of the enzymes triosephosphate isomerase (TIM) and alpha-glycerophosphate dehydrogenase (GDH) in a reaction mixture containing final concentrations of 100mM Tris pH 8.0, 4.75 mM fructose 1,6 bisphosphate, 0.15 mM NADH, 500 U/mL TIM, and 30 U/mL GDH.

The decrease in absorbance at 340nm, after addition of the cell extract supernatant, was recorded using an HP diode array spectrophotometer. One international unit (I.U.) of aldolase activity is that causing the oxidation of 2 µmol of NADH/min in this assay system.

Cell extracts containing the vector with the *fda* sequence showed a substantial increase in aldolase activity (13.1 I.U./mg protein) as compared to cells transformed with the control vector (0.15 I.U./mg protein). The activity was shown to be inhibited by EDTA, known to specifically inhibit class II aldolases.

EXAMPLE 2

Plant transformation and fda expression in tobacco

Targeting of FDA protein

E.coli fructose 1,6 bisphosphate aldolase was targeted to the plastid in plants in order to assess its influence on carbohydrate metabolism and starch biosynthesis in these plant organelles. To accomplish the import of the *E.coli* aldolase into the plastids, a vector was constructed in which the aldolase was fused to the *Arabidopsis* small subunit transit peptide (CTP1) (Stark et al., 1992) or the maize small subunit CTP (Russell et al., 1993), creating constructs in which the CTP-*fda* fusion gene was located between the 35S promoter from the figwort mosaic virus (P-FMV35S; Gowda et al., 1989) and the 3'-nontranslated region of the nopaline synthase gene (NOS 3'; Fraley et al., 1983) sequences. The vector construct containing the expression cassette [P-FMV/CTP1/*fda*/NOS3'] was subsequently used for tobacco protoplast transformation, which was performed as described in Fromm et al. (1987), with the following modifications. Tobacco cultivar Xanthi line D (Txd) cell suspensions were grown in 250-mL flasks, at 25°C and 138 rpm in the dark. For maintenance, a sub-culture volume of 9 mL was removed and added to 40 mL of fresh Txd media containing MS salts, 3% sucrose, 0.2 g/L inositol, 0.13 g/L asparagine, 80 µL of a 50 mg/mL stock of PCPA, 5 µL

of a 1 mg/mL stock of kinetin, and 1 mL of 1000x vitamins (1.3 g/L nicotinic acid, 0.25 g/L thiamine, 0.25 g/L pyridoxine HCL, and 0.25 g/L calcium pantothenate) every 3 to 4 days. Protoplasts were isolated from 1-day-old suspension cells that came from a 2-day-old culture. Sixteen milliliters of cells were added to 40 mL of fresh Txd media and
5 allowed to grow 24 hours prior to digestion and isolation of the protoplasts. The centrifugation stage for the enzyme mix has been eliminated. The electroporation buffer and protoplast isolation media were filter sterilized rather than autoclaved. The electroporation buffer did not have 4 mM CaCl_2 added. The suspension cells were digested in enzyme for 1 hour. Protoplasts were counted on a hemacytometer, counting
10 only the protoplasts that look intact and circular. Bio-rad Gene Pulser cuvettes (catalog # 165-2088) with a 0.4-cm gap and a maximum volume of 0.8 mL were used for the electroporations. Fifty to 100 μg of DNA containing the gene of interest along with 5 μg of internal control DNA containing the luciferase gene were added per cuvette. The final protoplast density at electroporation was $2 \times 10^6/\text{mL}$ and electroporator settings were a 500
15 μFarad capacitance and 140 volts on the Bio-rad Gene Pulser. Protoplasts were put on ice after resuspension in electroporation buffer and remained on ice in cuvettes until 10 minutes after electroporation. Protoplasts were added to 7 mL of Txd media + 0.4 M mannitol and conditioning media after electroporation. At this stage coconut water was no longer used. The protoplasts were grown in 1- hour day/night photoperiod regime at 26°C
20 and were spun down and assayed or frozen 20-24 hours after electroporation.

Western blot analysis performed on the protoplast extracts, obtained after transformation, showed processing into the mature FDA in the tobacco protoplasts. Expression was detected of a protein migrating at approximately 40 kDa, which is the molecular weight of the aldolase subunit and the size of the protein also observed after
25 overexpression of the aldolase in *E. coli*.

The expression cassette [P-FMV/CTP1/*fda*/NOS3'] was subsequently cloned into the NotI site of pMON17227 (described in PCT Publication WO 92/04449), in the same orientation as the selectable marker expression cassette, to form the plant transformation vector pMON17524, as shown in Figure 2 (SEQ ID NO: 5).

30 An additional construct was made and used for tobacco protoplast transformation, fusing the *fda* gene to the *Arabidopsis* EPSPS transit peptide (CTP2), which is described in US patent 5,463,175. The transit peptide was cloned (through the SphI site) into the

SphI site located immediately upstream from the N-terminus of the *fda* gene sequence in the CTP1-*fda* fusion (described above). This new CTP2-*fda* fusion gene was then cloned into a vector between the FMV promoter and the NOS 3' sequences. When this construct containing the CTP2/*fda* gene sequences was used for tobacco protoplast transformation, expression was detected of a protein migrating at approximately 40 kDa, which is the molecular weight of the aldolase subunit and the size of the protein also observed after overexpression of the aldolase in *E. coli*.

The NotI cassette [P-FMV/CTP2/*fda*/NOS3'] from this construct was then cloned into the NotI site of pMON17227, in the same orientation as the selectable marker expression cassette, to form the plant transformation vector pMON17542, which is shown in Figure 3 (SEQ ID NO:6).

For cytoplasmic expression of the FDA in tobacco protoplasts, a construct was made in which the *fda* gene sequence (without being coupled to a transit peptide) was cloned into a vector backbone, between the FMV promoter and the NOS 3' sequences. Using this construct for tobacco protoplast transformation also showed expression of a protein of the same size, migrating at approximately 40 kDa.

fda expression in tobacco plants

Two constructs, containing the *fda* gene, fused to the *Arabidopsis* small subunit CTP1 (pMON17524) (SEQ ID NO:5, Figure 2) and the *Arabidopsis* EPSPS (CTP2) transit peptide (pMON17542) (SEQ ID NO:6, Figure 3), were used for tobacco plant transformation, as described in US patent 5,463,175. A vector without the CTP-*fda* sequences, pMON17227 (described in PCT Publication WO 92/04449), was used as a negative control. The plant transformation vectors were mobilized into the ABI *Agrobacterium* strain. Mating of the plant vector into the ABI strain was done by the triparental conjugation system using the helper plasmid pRK2013 (Ditta et al., 1980).

Growth chamber-grown tobacco transformant lines were generated and first screened by Western blot analysis to identify expressors using goat antibody raised against *E.coli*-expressed *fda*. Subsequently, for pMON17524-expressing tobacco lines, leaf nonstructural carbohydrates were analyzed (sucrose, glucose, and hydrolyzed starch into glucose) by means of a YSI Instrument, Model 2700 Select Biochemistry Analyzer. Starting at flowering stage, leaf samples were also taken from these plants and analyzed for diurnal changes in leaf nonstructural carbohydrates.

Five hundred milligrams to 1 g fresh tobacco leaf tissue samples were harvested and extracted in 5 mL of hot Na-phosphate buffer (40 g/L NaH_2PO_4 and 10 g/L $\text{Na}_2\text{H}_2\text{PO}_4$ in double de-ionized water) by homogenization with a Polytron. Test tubes were then placed in an 85°C water bath for 15 minutes. Tubes were centrifuged for 12 minutes at 3000 rpm and the supernatants saved for soluble sugar analysis. The pellet was resuspended in 5 mL of hot Na-phosphate buffer mixed with a Vortex and centrifuged as described above. The supernatant was carefully removed and added to the previous supernatant fraction for soluble sugar (sucrose and glucose) analysis by YSI using appropriate membranes.

The starch was extracted from the pellet using the Megazyme Kit (Megazyme, Australia). To the pellet, 200 μL of 50% ethanol and 3 mL of thermostable alpha-amylase (300U) were added and the mixture vortexed. Samples were then incubated in boiling water for 6 minutes and stirred after 2 and 4 minutes. Tubes were placed in 50°C water bath and 4 mL of 200 mM acetate buffer (pH 4.5) were added followed by 0.1 mL amyloglucosidase (20 U). Incubation occurred for 1 hour. Test tubes were then centrifuged for 15 minutes at 3000 rpm. Aliquots were taken from the supernatant and analyzed for glucose by YSI. The free glucose was adjusted to anhydrous glucose (as it occurs in starch by multiplying by the ratio 162/182). The total volume per tube was 7.1 mL.

As seen in Table1, expression of the *fda* gene in tobacco correlated with a significant increase in leaf starch levels. However, referring to Figure 4, when a diurnal profile of starch levels was established in the *fda*-expressing leaves, this increase was apparent mainly early in the photoperiod, which is a phase when leaves are known to have peak photosynthetic activity. This increase in starch has no apparent negative effect on the plant because the increased starch is turned over during the dark period. There was no apparent increase in steady state levels of sucrose or glucose in tobacco leaves expressing *E.coli fda* as compared to the control.

Table 1
Leaf Carbohydrate Levels of Plants Expressing
the *fda* Transgene¹ (pMON17524)

5	High Expressors		Low Expressors	Negative
	(>0.01% total protein)		(< 0.01%)	Control
			(mg/g fresh weight)	
	STARCH	35.08 ± 2.84	23.25 ± 3.20	16.69 ± 2.92
10	SUCROSE	0.97 ± 0.17	0.86 ± 0.25	0.66 ± 0.19
	GLUCOSE	1.88 ± 0.17	1.58 ± 0.20	1.68 ± 0.26

¹ Leaf samples were harvested at midday.

A second set of transgenic tobacco plants transformed with the construct pMON17542 were grown in the greenhouse. Tobacco plants containing a vector without the CTP-*fda* sequences, pMON17227, were used as negative control. Of all the pMON17542-lines screened for expression by Western blot analysis, 18 were high expressors (>0.01% of the total cellular protein) and 15 lines were low expressors (<0.01%). Fifteen plants containing the null vector, pMON17227, were used as control.

Fully expanded leaves from plants expressing the *fda* transgene and negative controls were tested for sucrose export by collecting phloem exudate from excised leaf systems. The phloem exudation technique is described in Groussol et al. (1986). Leaves were harvested at 11:30 AM and placed in an exudation medium, containing 5 mM EDTA at pH 6.0, and allowed to exude for a period of 4 hours under full light and high humidity. The exudation solution was immediately analyzed for sucrose level, as described above in the carbohydrate analysis method. As seen in Table 2, a significant increase in sucrose export out of source leaves was observed in plants expressing the *fda* transgene.

This increase in sucrose export by *fda*-expressing leaves is an illustration of an increase in source capacity, very likely due to an increased carbon flow through the Calvin Cycle (in response to increased triose-P utilization) and thus an increase in net carbon utilization by the leaf. As seen in Table 2, the increase in sucrose loading in the phloem correlates with the level of *fda* expression.

Table 2
Levels of Sucrose in Phloem Exudate from
Excised Leaves of *fda* Transgenic Tobacco Plants (pMON17542)

	Water uptake (μ l/g F.Wt./h)	sucrose in phloem exudate	
		(ng/leaf)	(ng/g F.Wt.)
<i>fda</i> high expressors	320 \pm 20	330 \pm 60	108 \pm 22
<i>fda</i> low expressors	340 \pm 10	210 \pm 10	77 \pm 3
Control	390 \pm 30	160 \pm 10	56 \pm 3

Referring to Table 3, preliminary analysis of plant growth and development revealed no significant differences in number of leaves or pods per plant, plant height, stem diameter, or apparent seed weight per plant, between plants expressing the *fda* gene and the vector control under the specific growing and analysis conditions. However, as seen in Table 4, the *fda*-transgenic plants had a significantly higher root mass. This may be an indication that, under these conditions, roots represented a more dominant sink that attracted excess carbohydrate produced by the source leaves. Furthermore, the present illustration shows that the increase in root mass in the presence of the *E.coli fda* gene was accomplished with no apparent negative effect on shoot growth, inflorescence, or seed set. Therefore, this increase in root growth and final root dry weight is a desirable plant trait because it would lead to a rapid seedling establishment following germination and greater plant ability to tolerate drought, cold stress, other environmental challenges, and transplanting. In different plants and under different growing conditions, other plant parts (such as seed, fruit, stem, leaf, tuber, bulb, etc.) are expected to show the weight increase observed in tobacco roots overexpressing the *fda* transgene.

Table 3
Assessment of Certain Plant Growth and Development Parameters in
Tobacco Expressing the *fda* Transgene¹ (pMON17542)

	#pods/plant	#leaves/plant	Plant height (cm)	Seed weight (g/plant)
high expressors	162 ± 40	25.4 ± 0.8	65.3 ± 3.1	18.8 ± 2.4
Control	156 ± 28	24.4 ± 0.5	65.8 ± 5.1	17.3 ± 2.6

¹ To achieve this analysis, 14 high-expressor lines were compared to 15 control plants. Measurements were made prior to seed harvest (most pods have reached maturity). The number of leaves was confirmed by counting the number of nodes to account for leaf drop.

Table 4
Tobacco Root Dry Weight of Plants Expressing
the *E.coli fda* Transgene¹ (pMON17542)

	Root Dry Weight (g/plant)
<i>fda</i> high expressors	64.0 ± 3.9
<i>fda</i> low expressors	62.7 ± 5.4
Control	31.7 ± 1.6

¹ Roots from 5 high and 7 low expressing lines and 6 control plants were excised and washed carefully then placed in a 65°C drying oven for at least 48 hours. Roots were removed from the oven and allowed to equilibrate in the laboratory for 2 hours before dry weight determination.

EXAMPLE 3

Plant transformation and fda expression in corn plants

Targeting of FDA protein

Vectors containing the *fda* gene with and without the plastid targeting peptide were made for transformation in corn and are also suitable for other monocots, including rice, wheat, barley, sugarcane, triticale, etc.

For the cytosolic expression of the *fda* gene in corn plants, a construct was made in which the *fda* gene sequence was fused to the backbone of a vector containing the enhanced CaMV 35S promoter (e35S; Kay et al., 1987), the HSP70 intron (US patent 5,593,874), and the NOS3' polyadenylation sequence (Fraley et al., 1983). This created a NotI cassette [P-e35S/HSP70 intron/*fda*/NOS3'] that was cloned into the NotI site of pMON30460, a monocot transformation vector, to form the plant transformation vector pMON13925, as shown in Figure 5. pMON30460 contains an expression cassette for the selectable marker neomycin phosphotransferase typeII gene (nptII) [P-35S/NPTII /NOS3'] and a unique NotI site for cloning the gene of interest. The final vector (pMON13925) was constructed so that the gene of interest and the selectable marker gene were cloned in the same orientation. A vector fragment containing the expression cassettes for these gene sequences could be excised from the bacterial selector (Kan) and ori, gel purified, and used for plant transformation.

For the chloroplast-targeted expression of the *fda* gene in corn plants, a construct was made in which the *fda* gene sequence, coupled to the maize RUBISCO small subunit CTP (Russell et al., 1993), was fused to the backbone of a vector containing the enhanced (CaMV) 35S promoter, the HSP70 intron, and the NOS3' polyadenylation sequences. This created a NotI cassette [P-e35S/HSP70 intron/mzSSuCTP/*fda*/NOS3'] that was cloned into the NotI site (in the same orientation as the selectable marker cassette [P-35S/NPTII /NOS3']) of the monocot transformation vector pMON30460, to form the vector pMON17590, as shown in Figure 6. From this vector a fragment containing the *fda* gene expression cassette and the selectable marker cassette could be excised from the bacterial selector (Kan) and ori, gel purified, and used for plant transformation.

For the cytosolic endosperm-specific expression of the aldolase gene in corn, the *fda* gene sequence was cloned into a vector (in the same orientation as the selectable marker cassette[P-35S/NPTII /NOS3']) containing the glutelin gene promoter P-osgt1 (Zheng et al., 1993), the HSP70 intron, and the NOS3' polyadenylation sequences to form the vector pMON13936, as shown in Figure 7. From this vector a fragment containing the *fda* gene expression cassette [P-osgt1/HSP70intron/*fda*/NOS3'] and the selectable marker cassette could be excised from the bacterial selector (Kan) and ori, gel purified, and used for plant transformation.

Maize plant transformation

Transgenic maize plants transformed with the vectors pMON13925 (described above) or pMON17590 (described above) were produced using microprojectile bombardment, a procedure well-known to the art (Fromm, 1990; Gordon-Kamm et al., 1990; Walters et al., 1992). Embryogenic callus initiated from immature maize embryos was used as a target tissue. Plasmid DNA at 1 mg/mL in TE buffer was precipitated onto M10 tungsten particles using a calcium chloride / spermidine procedure, essentially as described by Klein et al. (1988). In addition to the gene of interest, the plasmids also contained the neomycin phosphotransferase II gene (nptII) driven by the 35S promoter from Cauliflower Mosaic Virus. The embryogenic callus target tissue was pretreated on culture medium osmotically buffered with 0.2M mannitol plus 0.2M sorbitol for approximately four hours prior to bombardment (Vain et al., 1993). Tissue was bombarded two times with the DNA-coated tungsten particles using the gunpowder version of the BioRad Particle Delivery System (PDS) 1000 device. Approximately 16 hours following bombardment, the tissue was subcultured onto a medium of the same composition except that it contained no mannitol or sorbitol, and it contained an appropriate aminoglycoside antibiotic, such as G418", to select for those cells that contained and expressed the 35S/nptII gene. Actively growing tissue sectors were transferred to fresh selective medium approximately every 3 weeks. About 3 months after bombardment, plants were regenerated from surviving embryogenic callus essentially as described by Duncan and Widholm (1988).

Aldolase activity from transgenic maize

In order to measure leaf aldolase activity, corn leaf samples were taken and immediately frozen on dry ice. Aldolase enzyme was extracted from the leaf tissue by grinding the leaf tissue at 4°C in 1.2 mL of the extraction buffer (100 mM Hepes, pH 8.0, 5 mM MgCl₂, 5 mM MnCl₂, 100 mM KCl, 10 mM DTT, 1% BSA, 1 mM PMSF, 10 µg/mL leupeptin, 10 µg/mL aprotinin). The extract was centrifuged at 15,000 x g, at 4°C for 3 minutes, and the non-desalted supernatant was assayed for enzyme activity. This extraction method gave about 60% recovery of *E. coli* FDA activity.

Total aldolase activity was determined in 0.98 mL of reaction mixture that consisted of 100 mM EPPS-NaOH, pH 8.5, 1 mM fructose-bisphosphate, 0.1 mM NADH, 5 mM MgCl₂, 4 units of alpha-glycerophosphate dehydrogenase, and 15 units of

triosephosphate isomerase. The reaction was initiated by addition of 20 μ L of leaf extract. The resulting data, generated from a single experiment, are presented in Table 5.

Table 5
Aldolase Activity from Transgenic Maize Leaves

5	Lines	A340/min/20 μ L	Activity %
	H99 (control)	0.113	100
	pMON 17590	0.233	206
	pMON13925	0.251	222

A phenotype was visible in the primary transformants (RO plants) expressing the *E. coli* FDA when the protein was targeted to the chloroplast. The leaves were chlorotic but seed set was normal. R1 plants were grown in both field and in greenhouse experiments. Starch was not detectable in the leaves using an iodine staining and pollination was delayed. It is believed that the phenotype in these corn plants may be the result of the promoter (e35S) used in both the pMON17590 and pMON13925 vectors not being preferred for causing FDA expression in corn. Because e35S is believed to cause mesophyll enhanced expression and the Calvin Cycle in a C4 plant such as corn occurs predominantly in the bundle sheath cells, the use of a promoter directing enhanced expression in the bundle sheath cells (such as the ssRUBISCO promoter) may be preferred. Vectors containing such a promoter and driving expression of FDA have been prepared and are being tested in maize.

In particular, the maize RuBISCO small subunit (PmzSSU, a bundle sheath cell-specific promoter) has been used to construct vectors for cell-specific *fda* expression in maize. A class I aldolase (*fdal*), an *fda* without an iron sulfur cluster and with different properties from *fdalI*, was utilized to improve carbon metabolism in C4 crops (e.g. maize). The gene for the class I aldolase was amplified from the genome of *Staphylococcus aureus* and activity was confirmed. Transformation vectors were then constructed to express both classes of aldolase (*fdal* and *fdalI*) in a cell-specific manner in maize. The following cassettes have been made:

pMON13899: PmzSSU/hsp70/mzSSU CTP/*fdal*
pMON13990: PmzSSU/hsp70/mzSSU CTP/*fdalI*

pMON13988:P35S/hsp70/*fda*.

These vectors were used for corn transformation as described generally above. The biochemical and physiological analysis of the primary transformants should allow for the identification of aldolase gene overexpression that will lead to increase growth and development and yield in maize.

Also, two vectors were used for transformation of corn which would target the expression of the *E. coli fda II* gene in the maize endosperm. The vector pMON 13936 uses the rice *gt1* promoter to drive expression of aldolase in the cytoplasm of the endosperm cells. Another vector (pMON 36416) uses the same promoter with the maize RuBISCO small subunit transit peptide to localize the protein in the amyloplasts. Homozygous lines of the cytosolic aldolase transformants have been identified (Homozygosity of 37 plants was confirmed using western blot analysis) and seed from these plants were collected for grain composition analysis (moisture, protein, starch, and oil). Of the 53 pMON 36416 primary transformants screened for amyloplast-targeted aldolase expression, 11 were positive. These plants will be tested for homozygosity selection/propagation and kernels from the homozygotes will be used for composition analysis.

EXAMPLE 4

Plant transformation and fda expression in potato plants

Targeting of fda expression

The plant expression vector, pMON17542 (described earlier), in which the *fda* gene is expressed behind the FMV promoter and the aldolase enzyme is fused to the chloroplast transit peptide CTP2, was used for *Agrobacterium*-mediated potato transformation.

A second potato transformation vector was constructed by cloning the NotI cassette [P-FMV/CTP2/*fda*/NOS3'] (described earlier) into the unique NotI site of pMON23616. pMON23616 is a potato transformation vector containing the nopaline-type T-DNA right border region (Fraley et al., 1985), an expression cassette for the neomycin phosphotransferase typeII gene [P-35S/NPTII /NOS3'] (selectable marker), a unique NotI site for cloning the gene expression cassette of interest, and the T-DNA left border region (Barker et al., 1983). Cloning of the NotI cassette [P-FMV/CTP2/*fda*/NOS3'] (described earlier) into the NotI site of pMON23616 results in the potato transformation vector

pMON17581, as shown in Figure 8. The vector pMON17581 was constructed such that the gene of interest and the selectable marker gene were transcribed in the same direction.

Potato plant transformation

The plant transformation vectors were mobilized into the ABI *Agrobacterium* strain. Mating of the plant vector into the ABI strain was done by the triparental conjugation system using the helper plasmid pRK2013 (Ditta et al., 1980). The vector pMON17542 was used for potato transformation via *Agrobacterium* transformation of Russet Burbank potato callus, following the method described in PCT Publication WO 96/03513 for glyphosate selection of transformed lines.

After transformation with the vector pMON17542, transgenic potato plantlets that came through selection on glyphosate were screened for expression of *E. coli* aldolase by leaf Western blot analysis. Out of 112 independent lines assayed, 50 *fda*-expressing lines (45%) were identified, with *fda* expression levels ranging between 0.12% and 1.2 % of total extractable protein.

The plant transformation vector PMON17581 was used for *Agrobacterium*-mediated transformation of HS31-638 potato callus. HS31-638 is a Russet Burbank potato line previously transformed with the mutant ADPglucose pyrophosphorylase (*glgC16*) gene from *E. coli* (U.S. Patent 5,498,830). The potato callus was transformed following the method described in PCT Publication WO 96/03513, substituting kanamycin (administered at a concentration of 150-200 mg/L) for glyphosate as a selective agent.

The transgenic potato plants were screened for expression of the *fda* gene by assaying leaf punches from tissue culture plantlets. Western blot analysis (using antibodies raised against the *E. coli* aldolase) of leaf tissue from the pMON17581-transformed lines identified 12 expressing lines out of 56 lines screened. Expression was detected of a protein migrating at approximately 40 kDa, which is the molecular weight of the *E. coli* (classII) aldolase subunit and the size of the protein observed after overexpression of the aldolase in *E. coli*.

Specific gravity measurements of transgenic potato plants

From the 50 *fda*-expressing potato lines obtained after transformation with pMON17542, 7 of the highest expressing lines were micropropagated in tissue culture, and 8 copies of each line were planted in pots 14 inches in diameter and 12 inches deep, containing a mixture of: ½ Metro 350 potting media, ¼ fine sand, ¼ Ready Earth

potting media. Wild-type Russet Burbank plantlets from tissue culture were planted as controls. All plants were cultivated for approximately 5 months in the greenhouse in which daytime temperature was approximately 21-23°C while nighttime temperature was approximately 13°C. Plants were watered every other day throughout their active growing
 5 period and fertilized with Peter's 20-20-20 commercial fertilizer once a week, at levels similar to commercial applications. Fertilization was carried out only for the first 2 ½ months, at which point fertilization was stopped completely. Plants were allowed to naturally senesce, and at approximately 50% senescence, tubers were harvested.

For each line at harvest, all tubers from all 8 pots were pooled and a total weight
 10 was obtained. Then for each line, tubers 30 g or greater were pooled and specific gravity was determined on this group of tubers. Specific gravity is the weight of the tubers in air divided by the weight in air minus the weight in water. Results of these weight measurements are presented in Table 6.

15 Table 6
 Specific gravity measurements from transgenic potato plants

Line #	Total Weight	Overall % Yield Increase	Combined Weight of Tubers over 30g	% Increase in Total Weight (Tubers over 30g)	Combined Weight of Tubers over 30g (% of Total Weight)	Specific Gravity
RB	6609		4477		67.70%	1.087
40652	5138	neg	1307	neg	25.40%	1.08
40611	7170	8.5%	4533	1.3%	63.20%	1.083
40608	7470	13.0%	1070	neg	14.30%	1.081
40632	7776	21.8%	5453	21.8%	70.10%	1.088
40614	8688	31.5%	5468	22.2%	62.90%	1.083
40631	8800	33.2%	6188	38.2%	70.30%	1.084
40610	9746	47.0%	7777	73.0%	80%	1.087

This table summarizes the tuber yield and specific gravity for all seven lines grown in the greenhouse. The results indicate that, in comparison to the control, all but one of the *fd*a lines show an increase in overall tuber yield, and that in four lines, there is a corresponding
 20 increase in percentage of tubers that weigh more than 30 g. For combined tubers over 30 g, the percent of total weight is near that of the control, and for two lines is greater than the control. This indicates that five out of the six of the lines show higher overall yield and are not making smaller tubers. In other words, with the increase in overall yield, there is a corresponding increase in percentage of bigger tubers (over 30 g). However, there is no
 25 increase in specific gravity of the tubers.

In conclusion, it appears that expression of *fda* in potato produces greater numbers of tubers per plant without a sacrifice in tuber size. This represents a yield benefit in that the farmer could potentially be able to produce the same amount of tubers using less acreage. Similar experiments will also be performed by co-expression of *fda* with other carbohydrate metabolizing genes, such as *glgC16*, in order to determine how such combinations will affect tuber yield, tuber solids deposition and overall tuber specific gravity.

Aldolase activity from transgenic potato

After being cultivated for 3 months (post planting) in the greenhouse, leaf samples were taken from 6 of the highest *fda*-expressing potato lines, obtained after transformation with pMON17542, and assayed for aldolase activity.

In order to measure potato leaf aldolase activity, duplicate leaf samples from each line were taken and immediately frozen on dry ice. Aldolase was extracted from 0.2 g of leaf tissue by grinding at 4°C in 1.2 mL of the extraction buffer: 100 mM Hepes, pH 8.0, 5 mM MgCl₂, 5 mM MnCl₂, 100 mM KCl, 10 mM DTT, 1% BSA, 1mM PMSF, 10 µg/mL leupeptin, 10 µg/mL aprotinin. The extract was assayed for aldolase activity as described earlier.

Six independent transgenic potato lines expressing *fda* were tested for aldolase activity. The expression of *fda* in leaves is an indicator of the expression in the whole plant because the FMV promoter used to drive expression of the respective encoding DNAs directs gene expression constitutively in most, if not all, tissues of potato plants.

Table 7 summarizes the quantitative protein expression data for each of the lines, and the percent activity for each individual line.

Table 7
Aldolase Activity from
Transgenic Russet Burbank Potato Leaves

5	Lines	Exp. #1		Exp. #2		Average
		Act (U/gFW)	%Act	Act (U/gFW)	%Act	% Activity
	Control	4.461	100	4.732	100	100
	40608	6.969	156	8.055	170	163
	40610	8.489	190	7.326	155	173
	40652	5.812	130	6.367	135	132
10	40632	5.257	118	4.244	90	104
	40631	5.764	129	4.968	105	117
	40611	5.715	128	5.836	123	126

Solids uniformity in transgenic potato

15 Twenty-five Russet Burbank lines expressing *fda* (potato lines designated "Maestro"), obtained after transformation with pMON17542, and fifteen Russet Burbank Simple Solid lines, also containing *glgC16* (PCT Publication WO 91/19806 and US Patent 5,498,830), expressing *fda* (potato lines designated "Segal"), obtained after transformation with pMON17581, were field tested at two different sites. For each field site, 36 plants
20 per line (three repetitions of 12 plants per line) were evaluated for tuber solids distribution. At harvest, tubers were pre-sorted at each field site into a ten to twelve ounce category, and nine tubers from each replicated plot were analyzed in groups of three.

For a typical 10-12 ounce tuber having a diameter of 7-8 cm, starch distribution was evaluated by removing the center longitudinal slice (13 mm) from each tuber. Slices
25 were then peeled and laid flat on a cutting board where the inner tuber region (pith region) was removed by a 14-mm cork punch. The tissue from pith to cortex (perimedullary region) was removed by an up-to-a 2-inch cork punch. The remaining cortex tissue was approximately an 8-mm wide ring from the outermost region of the slice.

Specific gravity was then determined by weighing both the pooled pith punches
30 and pooled cortex punches in air and then in water:

$$\text{Specific gravity} = \text{Air Wt.}/(\text{Air Wt.}-\text{Water Wt.})$$

After calculating specific gravity, solids levels were determined by the following equation:

$$-214.9206 + (218.1852 * \text{Sp. Gravity})$$

The degree of solids uniformity (Solids Uniformity Index) is determined by calculating the pith to cortex solids ratio (pith solids divided by cortex solids). The three groups of three
5 tubers per plot were averaged, at which point the average of three plot replications was calculated per field site.

Analyses of several previous solids uniformity field trials (data not shown) have demonstrated nontransgenic, wild-type Russet Burbank potato to have a typical pith to cortex tuber solids ratio within the range of 68% to 72%, depending on growing region
10 and agricultural practices. Tables 8-11 provide the pith to cortex solids ratios by plant line number, with a higher pith to cortex solids ratio indicating a greater degree of solids uniformity.

Tables 8 and 9 represent the data from one field site (site 1) for Segal and Maestro, respectively, and illustrate that the majority of Segal and Maestro lines have higher pith to
15 cortex solids ratios than that of 68.4% for the Russet Burbank control, with some lines approaching an 82% pith to cortex solids ratio.

Tables 10 and 11 represent the data from another field site (site 2) for Segal and Maestro, respectively, and also illustrate that the majority of Maestro and Segal lines have higher pith to cortex solids ratios than that of the Russet Burbank control, with some lines
20 approaching an 88% pith to cortex solids ratio. In the site 2 field trial, the Russet Burbank control had an atypical, abnormally high pith-to-cortex solids uniformity ratio of 79.3%, which was most likely due to environmental growing conditions. The site 2 results demonstrate that expression in Russet Burbank potato of *E. coli fda*, alone or with co-expression of *glgC16*, increases tuber solids uniformity even in a growing season when
25 tuber solids uniformity is already extremely high in nontransgenic Russet Burbank. That is, the *fda* gene continues to perform when agricultural conditions are already conducive to an abnormally high solids uniformity level.

**Table 8. Solids Uniformity Index: Pith Solids to Cortex Solids Ratio.
Segal Russet Burbank Lines. Site 1**

	<u>Line</u>	<u>Ratio</u>
5	S-29	79.1
	S-9	75.8
	S-20	71.3
	S-15	71.3
	S-21	70.5
10	S-5	70.2
	S-18	70.0
	RB control	68.4
	S-32	68.3
	S-16	65.6
15		

**Table 9. Solids Uniformity Index: Pith Solids to Cortex Solids Ratio.
Maestro Russet Burbank Lines. Site 1**

	<u>Line</u>	<u>Ratio</u>
20	M-13	74.0
	M-12	73.6
	M-1	73.4
	M-3	73.0
	M-6	72.4
25	M-9	71.2
	M-11	70.6
	M-18	70.5
	M-17	69.9
	M-19	69.4
30	M-5	69.3
	M-20	68.9
	RB control	68.4
	M-8	68.3
	M-43	67.7
35	M-23	67.3
	M-7	67.0
	M-39	66.6
	M-22	66.0
	M-10	65.4
40	M-27	61.4

**Table 10. Solids Uniformity Index: Pith Solids to Cortex Solids Ratio
Segal Russet Burbank Lines. Site 2**

	<u>Line</u>	<u>Ratio</u>
5	S-33	87.4
	S-54	87.1
	S-05	86.8
	S-29	85.1
	S-21	84.3
10	S-16	83.2
	S-20	81.5
	S-18	80.7
	S-32	80.6
	RB control	79.3
15	S-09	79.0

**Table 11. Solids Uniformity Index: Pith Solids to Cortex Solids Ratio
Maestro Russet Burbank Lines. Site 2**

	<u>Line</u>	<u>Ratio</u>
20	M-04	87.7
	M-18	83.9
	M-17	83.8
	M-03	83.7
	M-09	83.4
25	M-15	83.2
	M-29	82.9
	M-44	82.3
	M-08	82.2
	M-43	81.6
30	M-22	81.1
	M-05	80.8
	M-01	80.5
	M-20	80.2
	M-45	79.6
35	M-39	79.5
	M-27	79.5
	RB control	79.3
	M-13	78.9
	M-22	78.8
40	M-19	78.7
	M-07	78.2
	M-12	77.9
	M-23	77.3
	M-06	76.5
45	M-10	75.0
	M-11	74.1

The effect of aldolase on pith to cortex solids ratios in the Segal lines is slightly more dramatic than in Maestro lines. We believe this phenotype is due to expression of *fda* in a background in which the Russet Burbank host expresses *glgC16* at a relatively low to moderate level, and that the combination of *fda* plus *glgC16* provides improved
5 benefits. Cross sectional tuber slices (Figure 9) of three Segal lines with improved solids uniformity illustrate a greater deposition of starch within the inner regions of the tuber. Specifically, an increase in cortex volume accompanied by relocation of the xylem ring towards the center of the tuber, plus a more opaque pith tissue due to an increase in starch density, are evident in the transgenic lines. This physiological alteration may be due to an
10 increase in sucrose translocation from source to sink, which may influence phloem element distribution during tuber development or sucrose availability for starch biosynthesis across the tuber.

Example 5

Plant transformation and FDA expression in cotton plants

15 The *E. coli fda* vectors pMON17524 [FMV/CTP1/*fda*] (Figure 2) and pMON17542 [FMV/CTP2/*fda*] (Figure 3) were transformed into cotton using *Agrobacterium* as described by Umbeck et al. (1987) and in US Patent 5004863. The protein was targeted to the chloroplast using either the *Arabidopsis* SSU CTP 1 (pMON17524) or the *Arabidopsis* EPSPS (pMON17542) chloroplast transit peptide.

20 *Aldolase expression in cotton*

Five-week-old calli transformed with both vectors were analyzed by Western blot analyses and by aldolase assays. Western blot analysis indicated a large amount of protein at the position of the full-length FDA standard and a lesser amount at the same position in the control callus extracts. It appeared that the protein was fully processed. To verify that
25 FDA was expressed in the tissue and for comparison of activity, calli transformed with the two vectors were extracted in a buffer that would prevent loss of activity of the transgene product. BSA was added to final concentration of 1 mg/mL, which limited the analysis of processing on import by Western blot. Aldolase assays were performed plus or minus 25 mM EDTA, which inhibits the *E. coli* enzyme but not the plant enzyme. The results of the
30 assays are shown in Table 12.

Table 12

Aldolase Activity in Cotton Calli and Cotton Leaf

 $\Delta A_{340} \text{ e}^{-3}/\text{mg protein}/5 \text{ min}$

	Colony#	-EDTA	+EDTA	Fold Increase
5	<u>Controls</u>			
	Cotton Leaf (Coker)	4.0	4.2	-
	Uninoculated Calli	7.7	5.6	1.3X
	Inoculated Calli (E35S/GUS) #1	6.8	6.1	-
	#2	3.5	4.0	-
10	<u>FDA calli</u>			
	pMON 17542 #1	3.5	2.3	1.5X
	#3	5.5	2.6	2.1X
	#5	9.2	3.8	2.4X
	#4	19.8	3.6	5.5X
15	pMON17524 #2	15.2	5.8	2.6X
	#3	12.5	4.0	3.1X
	#5	14.4	2.9	4.9X
	#6	4.1	1.2	3.5X

- 20 The results indicate that there is good expression of the *fda* gene in cotton callus. Almost all calli had at least twofold higher aldolase activity, and the increase was sensitive to inhibition by EDTA. Processing appeared complete by Western blot analysis using these samples.

25

REFERENCES CITED

- Alefunder et al. (1989) Biochem. J. 257:529-534
- Ammirato et al. (1984) Handbook of Plant Cell Culture - Crop Species. Macmillan Publ.
 5 Co..
- Bai et al. (1975) Arch. Biochem. Biophys. 168: 230-234.
- Baldwin et al. (1978) Biochem. J. 169: 633-641
- Barker et al. (1983) Plant Mol Biol 2 (6): 335-350.
- Benfey et al. (1989) EMBO J. 5: 2195-2202.
- 10 Besmond et al. (1983) Biochem. Biophys. Res. Commun. 117, 601-609.
- Bevan (1984) Nucleic Acids Res. 12 (22): 8711-8721.
- Bevan et al. (1986) Nucleic Acids Res. 14 (11):4625-4638.
- Campbell et al. (1994) Canadian Journal of Forest Research 24 (8):1689-1693.
- Cerdan et al. (1997) Plant Molecular Biology 33 (2): p245-255.
- 15 Chopra et al. (1990) Plant Molecular Biology 15:517-520.
- Clayton (1985) EMBO J. 4, 2997-3003.
- Cremona (1968) G. Bot. Ital. 102, 253-259.
- Daniell et al. (1998) Nature Biotechnology 16:345-348.
- Datta et al. (1990) Bio-technology 8:736-740.
- 20 Ditta et al. (1980) Proc Natl Acad Sci USA 77(12): 7347-7351.
- Duncan and Widholm (1988) Plant Cell Reports 7: 452-455.
- Edwards et al. (1990). Proc Natl Acad Sci USA 87 (9): p3459-3463.
- Fejes et al. (1990). Plant Mol Biol 15 (6): p921-932.
- Fraley et al. (1983) Proc Natl Acad Sci USA 80: 4803-4807.
- 25 Fraley et al. (1985) Bio/Technology 3 (7): 629-635.
- Fromm, M., (1990) UCLA Symposium on Molecular Strategies for Crop Improvement,
 April 16-22, 1990. Keystone, CO.
- Fromm et al. (1987) Methods in Enzymology. 153:351-366.
- Gordon-Kamm et al. (1990) Plant Cell 2: 603-618.
- 30 Gotz et al. (1979) FEMS Microbiol. Lett. 5:253-257.
- Gowda et al. (1989). Journal of Cellular Biochemistry supplement 13D, 301 (Abstract).
- Groussol et al. (1986) Physiologie Vegetale 24(1):123-134.

- Guerrini et al. (1971) Arch. Biochem. Biophys. 146, 249-255.
- Hannapel (1990) Plant Physiol. 94: 919-925.
- Hayashimoto et al. (1990) Plant Physiol. 93:857-863.
- Herrera-Estrella et al. (1983) Nature 303:209.
- 5 Hinchee et al., Bio/Technology 6:915-922 (1988).
- Horsch and Klee. (1986) Proc. Natl. Acad. Sci. U.S.A. 83:4428-4432.
- Jack (1973) Ph.D. Dissertation, University of Cambridge.
- Jack and Harris (1971) Biochem. J. 124, 680-690.
- Jefferson et al. (1990) Plant Mol. Biol. 14: 995-1006.
- 10 Joh et al. (1986) J. Mol. Biol. 190:401-410.
- Kay et al. (1987) Science 236: 1299-1302.
- Klee et al. (1985) Bio-Technology 3(7): 637-642.
- Klein et al. (1988) Bio/Technology 6: 559-563.
- Kretsch et al. (1995) Plant Journal 7 (5): p715-729.
- 15 Lai et al., (1974) Science 183, 1204-1206.
- Lebherz and Rutter (1973) J. Biol. Chem. 248:1650-1659.
- Lebherz et al (1984) J. Biol. Chem. 259 (2):1011-1017.
- Leyva et al. (1995) Plant Physiology 108(1):39-46.
- London and Kline (1973) Bacteriol. Rev. 37:453-478.
- 20 Lloyd et al. (1991). Mol. Gen. Genet. 225 (2):209-216.
- Luan et al. (1992). Plant Cell 4 (8):971-981.
- Luebberstedt et al. (1994) Plant Physiology 104 (3):997-1006.
- Maniatis et al. (1982) Molecular Cloning: A laboratory manual. Cold Spring Harbor
Laboratory Press, Cold Spring Harbor, N.Y.
- 25 Matsuoka et al. (1993). Proc. Natl. Acad. Sci. U. S. A. 90(20):9586-9590.
- Mignery et al (1988) Gene 62:27-44.
- Muller et al (1990) Mol. Gen. Genet. 224:136-146.
- Oelmueller et al. (1992). Res. Photosynth., Proc. Int. Congr. Photosynth., 9th.
Volume 3: p219-24. Editor(s): Murata, Norio. Publisher: Kluwer, Dordrecht,
- 30 Neth.
- Pedersen et al. (1982) Cell 29:1015-1026.
- Potrykus et al. (1985), Mol. Gen. Genet. 199:183-188.

- Rocha-Sosa et al. (1989) EMBO J. 8 (1):23-29.
- Rogers et al. (1987) Improved vectors for plant transformation: expression cassette vectors and new selectable markers. In Methods in Enzymology. Edited by R. Wu and L. Grossman. p253-277. San Diego: Academic Press.
- 5 Rohde et al. (1990) J. Genet. & Breed. 44:311-315.
- Russell et al. (1993) Plant Cell Reports 13:24-27.
- Russell and Fromm (1997) Transgenic Research 6 (2):157-168.
- Russel and Gibbs (1967) Biochim. Biophys. Acta 132, 145-154
- Salanoubat and Belliard (1987) Gene 60:47-56.
- 10 Salanoubat and Belliard (1989) Gene 84:181-185.
- Samac et al. (1990) Plant Physiol. 93:907-914.
- Sambrook et al. (1989) Molecular cloning: A laboratory manual, 2nd ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N. Y.
- Schmidhauser and Helinski. (1985) J. Bacteriol. 164-155.
- 15 Sonnewald et al. (1994) Plant Cell and Environment 17:649-658.
- Stalker et al. (1988) J. Biol. Chem. 263:6310-6314.
- Stark et al. (1992) Science 258: 287-292.
- Stockhaus et al. (1989). EMBO Journal 8(9):2445-2451.
- Stribling and Perham (1973) Biochem. J. 131:833-841.
- 20 Suzuki et al. (1994) Plant Mol. Biol. 25(3):507-516.
- Thillet et al. (1988) J. Biol. Chem. 263:12500-12508.
- Thompson et al. (1994) Nucl. Acids Res. 22:4673-4680.
- Tierney et al. (1987) Planta 172:356-363.
- Truernit et al. (1995) Planta 196 (3):564-570.
- 25 Tsutsumi et al. (1984) J. Biol. Chem. 259, 14572-14575.
- Umbeck et al. (1987) Biotechnology. 5, 263-266.
- Vain et al. (1993) Plant Cell Reports 12: 84-88.
- Vasil et al. (1990) Bio/Technology 8:429-434.
- Vasil et al. (1992) Bio/Technology 10:667-674.
- 30 Walters et al. (1992) Plant Molecular Biology 18: 189-200.
- Witke and Goetz (1993) Journal of Bacteriology 175(22): 7495-7499
- Wong et al. (1988) Gene 68: 193-203.

Yamamoto et al. (1994) Plant and Cell Physiology 35(5):773-778.

Zheng et al. (1993) Plant J. 4: 3357-3366.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

5

(i) APPLICANT: Gerard Barry
NordineCheikh
Ganesh Kishore

10

(ii) TITLE OF INVENTION: Expression of Fructose 1,6 Bisphosphate
Aldolase in Transgenic Plants

(iii) NUMBER OF SEQUENCES: 6

15

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Arnold, White & Durkee
(B) STREET: P.O. Box 4433
(C) CITY: Houston
(D) STATE: Texas
(E) COUNTRY: United States of America
(F) ZIP: 77210-4433

20

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30

25

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER: US Unknown
(B) FILING DATE: Concurrently Herewith
(C) CLASSIFICATION: Unknown

30

(vi) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: US Prov. App. Serial No. 60/049,995
(B) FILING DATE: June 17, 1997

35

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Patricia A. Kammerer
(B) REGISTRATION NUMBER: 29,775
(C) REFERENCE/DOCKET NUMBER: MOBT086

40

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (713) 787-1400
(B) TELEFAX: (713) 787-1440

45

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1080 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single

55

(D) TOPOLOGY: linear

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

10 ATGTCTAAGA TTTTGTGATT CGTAAAACCT GCGTAATCA CTGGTGATGA CGTACAGAAA
60

GTTTTCCAGG TAGCAAAAGA AAACAACCTC GCACTGCCAG CAGTAACTG CGTCGGTACT
120

15 GACTCCATCA ACGCCGTACT GGAAACCGCT GCTAAAGTTA AAGCGCCGGT TATCGTTCAG
180

TTCTCCAACG GTGGTGCTTC CTTTATCGCT GGTAAGGCG TGAAATCTGA CGTTCCGCAG
240

20 GGTGCTGCTA TCCTGGGCGC GATCTCTGGT GCGCATCACG TTCACCAGAT GGCTGAACAT
300

TATGGTGTTC CGGTTATCCT GCACACTGAC CACTGCGCGA AGAACTGCT GCCGTGGATC
25 360

GACGGTCTGT TGGACGCGGG TGA AAAACAC TTCGCAGCTA CCGGTAAGCC GCTGTTCTCT
420

30 TCTCACATGA TCGACCTGTC TGAAGAATCT CTGCAAGAGA ACATCGAAAT CTGCTCTAAA
480

TACCTGGAGC GCATGTCCAA AATCGGCATG ACTCTGGA AA TCGAACTGGG TTGCACCGGT
540

35 GGTGAAGAAG ACGGCGTGGA CAACAGCCAC ATGGACGCTT CTGCACTGTA CACCCAGCCG
600

GAAGACGTTG ATTACGCATA CACCGAACTG AGCAAAATCA GCCCGCGTTT CACCATCGCA
40 660

GCGTCCTTCG GTAACGTACA CGGTGTTTAC AAGCCGGGTA ACGTGGTTCT GACTCCGACC
720

45 ATCCTGCGTG ATTCTCAGGA ATATGTTTCC AAGAAACACA ACCTGCCGCA CAACAGCCTG
780

AACTTCGTAT TCCACGGTGG TTCCGGTTCT ACTGCTCAGG AAATCAAAGA CTCCGTAAGC
840

50 TACGGCGTAG TAAAAATGAA CATCGATACC GATACCCAAT GGGCAACCTG GGAAGGCGTT
900

CTGAACTACT ACAAAGCGAA CGAAGCTTAT CTGCAGGGTC AGCTGGGTAA CCCGAAAGGC
55 960

GAAGATCAGC CGAACAAGAA ATACTACGAT CCGCGCGTAT GGCTGCGTGC CGGTCAGACT
1020

TCGATGATCG CTCGTCTGGA GAAAGCATTC CAGGAACTGA ACGCGATCGA CGTTCTGTAA
1080

5

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- 10 (A) LENGTH: 359amino acids
(B) TYPE: amino
(C) STRANDEDNESS:
(D) TOPOLOGY: Linear

15 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2

	Met	Ser	Lys	Ile	Phe	Asp	Phe	Val	Lys	Pro	Gly	Val	Ile	Thr	Gly	
				5					10						15	
20	Asp	Asp	Val	Gln	Lys	Val	Phe	Gln	Val	Ala	Lys	Glu	Asn	Asn	Phe	
				20					25						30	
	Ala	Leu	Pro	Ala	Val	Asn	Cys	Val	Gly	Thr	Asp	Ser	Ile	Asn	Ala	
25				35					40						45	
	Val	Leu	Glu	Thr	Ala	Ala	Lys	Val	Lys	Ala	Pro	Val	Ile	Val	Gln	
				50					55						60	
	Phe	Ser	Asn	Gly	Gly	Ala	Ser	Phe	Ile	Ala	Gly	Lys	Gly	Val	Lys	
30				65					70						75	
	Ser	Asp	Val	Pro	Gln	Gly	Ala	Ala	Ile	Leu	Gly	Ala	Ile	Ser	Gly	
				80					85						90	
35	Ala	His	His	Val	His	Gln	Met	Ala	Glu	His	Tyr	Gly	Val	Pro	Val	
				95					100						105	
	Ile	Leu	His	Thr	Asp	His	Cys	Ala	Lys	Lys	Leu	Leu	Pro	Trp	Ile	
40				110					115						120	
	Asp	Gly	Leu	Leu	Asp	Ala	Gly	Glu	Lys	His	Phe	Ala	Ala	Thr	Gly	
				125					120						135	
	Lys	Pro	Leu	Phe	Ser	Ser	His	Met	Ile	Asp	Leu	Ser	Glu	Glu	Ser	
45				140					145						150	
	Leu	Gln	Glu	Asn	Ile	Glu	Ile	Cys	Ser	Lys	Tyr	Leu	Glu	Arg	Met	
				155					160						165	
50	Ser	Lys	Ile	Gly	Met	Thr	Leu	Glu	Ile	Glu	Leu	Gly	Cys	Thr	Gly	
				170					175						180	
	Gly	Glu	Glu	Asp	Gly	Val	Asp	Asn	Ser	His	Met	Asp	Ala	Ser	Ala	
				185					190						195	
55	Leu	Tyr	Thr	Gln	Pro	Glu	Asp	Val	Asp	Tyr	Ala	Tyr	Thr	Glu	Leu	
				200					205						210	

	Ser Lys Ile Ser Pro Arg Phe Thr Ile Ala Ala Ser Phe Gly Asn	
	215	220 225
5	Val His Gly Val Tyr Lys Pro Gly Asn Val Val Leu Thr Pro Thr	
	230	235 240
	Ile Leu Arg Asp Ser Gln Glu Tyr Val Ser Lys Lys His Asn Leu	
	245	250 255
10	Pro His Asn Ser Leu Asn Phe Val Phe His Gly Gly Ser Gly Ser	
	260	265 270
	Thr Ala Gln Glu Ile Lys Asp Ser Val Ser Tyr Gly Val Val Lys	
	275	280 285
15	Met Asn Ile Asp Thr Asp Thr Gln Trp Ala Thr Trp Glu Gly Val	
	290	295 300
	Leu Asn Tyr Tyr Lys Ala Asn Glu Ala Tyr Leu Gln Gly Gln Leu	
20	305	310 315
	Gly Asn Pro Lys Gly Glu Asp Gln Pro Asn Lys Lys Tyr Tyr Asp	
	320	325 330
25	Pro Arg Val Trp Leu Arg Ala Gly Gln Thr Ser Met Ile Ala Arg	
	335	340 345
	Leu Glu Lys Ala Phe Gln Glu Leu Asn Ala Ile Asp Val Leu	
	350	355
30		

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

35 (A) LENGTH: 30 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: Linear

40 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

GGGGCCATGG CTAAGATTTT TGATTTCGTA

(2) INFORMATION FOR SEQ ID NO:4:

45 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 34 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

50 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

CCCCGAGCTC TTACAGAACG TCGATCGCGT TCAG

55

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 10847 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: Linear

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

```

      1  CGATAAGCTT  GATGTAATTG  GAGGAAGATC  AAAATTTTCA  ATCCCCATTC
    51  TTCGATTGCT  TCAATTGAAG  TTTCTCCGAT  GCGCAAGTT  AGCAGAATCT
  10  101  GCAATGGTGT  GCAGAACCCA  TCTCTTATCT  CCAATCTCTC  GAAATCCAGT
    151  CAACGCAAT  CTCCCTTATC  GGTTCCTCTG  AAGACGCAGC  AGCATCCACG
    201  AGCTTATCCG  ATTTTCGTCT  CGTGGGGATT  GAAGAAGAGT  GGGATGACGT
    251  TAATTGGCTC  TGAGCTTCGT  CCTCTTAAGG  TCATGTCTTC  TGTTTCCACG
    301  GCGTGCATGC  TTCACGGTGC  AAGCAGCCGT  CCAGCAACTG  CTCGTAAGTC
  15  351  CTCTGGTCTT  TCTGGAACCG  TCCGTATTCC  AGGTGACAAG  TCTATCTCCC
    401  ACAGGTCCTT  CATGTTTGGA  GGTCTCGCTA  GCGGTGAAAC  TCGTATCACC
    451  GGTCTTTTGG  AAGGTGAAGA  TGTTATCAAC  ACTGGTAAGG  CTATGCAAGC
    501  TATGGGTGCC  AGAATCCGTA  AGGAAGGTGA  TACTTGGATC  ATTGATGGTG
    551  TTGGTAACGG  TGGACTCCTT  GCTCCTGAGG  CTCCTCTCGA  TTTGCGTAAC
  20  601  GCTGCAACTG  GTTGCCGTTT  GACTATGGGT  CTTGTTGGTG  TTTACGATTT
    651  CGATAGCACT  TTCATTGGTG  ACGCTTCTCT  CACTAAGCGT  CCAATGGGTC
    701  GTGTGTTGAA  CCCACTTCGC  GAAATGGGTG  TGCAGGTGAA  GTCTGAAGAC
    751  GGTGATCGTC  TTCCAGTTAC  CTTGCGTGGA  CCAAAGACTC  CAACGCCAAT
    801  CACCTACAGG  GTACCTATGG  CTTCCGCTCA  AGTGAAGTCC  GCTGTTCTGC
  25  851  TTGCTGGTCT  CAACACCCCA  GGTATCACCA  CTGTTATCGA  GCCAATCATG
    901  ACTCGTGACC  ACACTGAAAA  GATGCTTCAA  GGTTCCTGAT  CTAACCTTAC
    951  CGTTGAGACT  GATGCTGACG  GTGTGCGTAC  CATCCGTCTT  GAAGGTCGTG
  1001  GTAAGCTCAC  CGGTCAAGTG  ATTGATGTTT  CAGGTGATCC  ATCCTCTACT
  1051  GCTTTCCCAT  TGGTTGCTGC  CTTGCTTGTT  CCAGGTTCCG  ACGTCACCAT
  30  1101  CCTTAACGTT  TTGATGAACC  CAACCCGTAC  TGGTCTCATC  TTGACTCTGC
    1151  AGGAAATGGG  TGCCGACATC  GAAGTGATCA  ACCCACGTCT  TGCTGGTGGA
    1201  GAAGACGTGG  CTGACTTGCG  TGTTGCTTCT  TCTACTTTGA  AGGGTGTTAC
    1251  TGTTCAGAA  GACCGTGCTC  CTTCTATGAT  CGACGAGTAT  CCAATTCTCG
    1301  CTGTTGAGC  TGCATTGCT  GAAGGTGCTA  CCGTTATGAA  CGGTTTGAA
  35  1351  GAACTCCGTG  TTAAGGAAAG  CGACCGTCTT  TCTGCTGTCG  CAAACGGTCT
    1401  CAAGCTCAAC  GGTGTTGATT  GCGATGAAGG  TGAGACTTCT  CTCGTCGTGC
    1451  GTGGTCGTCC  TGACGGTAAG  GGTCTCGGTA  ACGCTTCTGG  AGCAGCTGTC
    1501  GCTACCCACC  TCGATCACCG  TATCGCTATG  AGCTTCCTCG  TTATGGGTCT
    1551  CGTTTCTGAA  AACCTGTGTA  CTGTTGATGA  TGCTACTATG  ATCGCTACTA
  40  1601  GCTTCCCAGA  GTTCATGGAT  TTGATGGCTG  GTCTTGGAGC  TAAGATCGAA
    1651  CTCTCCGACA  CTAAGGCTGC  TTGATGAGCT  CAAGAATTCT  AGCTCGGTAC
    1701  CGGATCCAGC  TTTCGTTCTG  ATCATCGGTT  TCGACAACGT  TCGTCAAGTT
    1751  CAATGCATCA  GTTCATTGCT  GCACACACCA  GAATCCTACT  GAGTTCGAGT
    1801  ATTATGGCAT  TGGGAAAAC  GTTTTCTTGG  TACCATTTGT  TGTGCTTGTA
  45  1851  ATTTACTGTG  TTTTATATTC  GGTTCGCT  ATCGAACTGT  GAAATGGAAA
    1901  TGGATGGAGA  AGAGTTAATG  AATGATATGG  TCCTTTTGTT  CATTCTCAA
    1951  TTAATATTAT  TTGTTTTTTC  TCTTATTTGT  TGTGTGTTGA  ATTTGAAATT
    2001  ATAAGAGATA  TGCAAACATT  TTGTTTGTAG  TAAAAATGTG  TCAAATCGTG
    2051  GCCTCTAATG  ACCGAAGTTA  ATATGAGGAG  TAAACACTT  GTAGTTGTAC
  50  2101  CATTATGCTT  ATTCACTAGG  CAACAAATAT  ATTTTCAGAC  CTAGAAAAGC
    2151  TGCAAATGTT  ACTGAATACA  AGTATGTCCT  CTTGTGTTT  AGACATTTAT
    2201  GAACTTTCTT  TTATGTAATT  TTCCAGAATC  CTTGTGAGAT  TCTAATCATT
    2251  GCTTTATAAT  TATAGTTATA  CTCATGGATT  TGTAGTTGAG  TATGAAAATA
    2301  TTTTATAATG  CATTATATGA  CTTGCCAATT  GATTGACAAC  ATGCATCAAT
  55  2351  CGACCTGCAG  CCACTCGAAG  CGGCCGCGTT  CAAGCTTGAG  CTCAGGATTT
    2401  AGCAGCATTC  CAGATTGGGT  TCAATCAACA  AGGTACGAGC  CATATCACTT
    2451  TATTCAAATT  GGTATCGCCA  AAACCAAGAA  GGAACCTCCA  TCCTCAAAGG
    2501  TTTGTAAGGA  AGAATTCTCA  GTCCAAAGCC  TCAACAAGGT  CAGGGTACAG

```

2551 AGTCTCCAAA CCATTAGCCA AAAGCTACAG GAGATCAATG AAGAATCTTC
 2601 AATCAAAGTA AACTACTGTT CCAGCACATG CATCATGGTC AGTAAGTTTC
 2651 AGAAAAAGAC ATCCACCGAA GACTTAAAGT TAGTGGGCAT CTTTGAAAGT
 2701 AATCTTGTCA ACATCGAGCA GCTGGCTTGT GGGGACCAGA CAAAAAGGA
 5 2751 ATGGTGCAGA ATTGTTAGGC GCACCTACCA AAAGCATCTT TGCCTTTATT
 2801 GCAAAGATAA AGCAGATTCC TCTAGTACAA GTGGGGAACA AAATAACGTG
 2851 GAAAAGAGCT GTCCTGACAG CCCACTCACT AATGCGTATG ACGAACGCAG
 2901 TGACGACCAC AAAAGAATTC CCTCTATATA AGAAGGCATT CATTCCCATT
 2951 TGAAGGATCA TCAGATACTG AACCAATCCT TCTAGAAGAT CTCCACAATG
 10 3001 GCTTCTCTTA TGCTCTCTTC CGCTACTATG GTTGCCTCTC CGGCTCAGGC
 3051 CACTATGGTC GCTCCTTTCA ACGGACTTAA GTCTCCGCT GCCTTCCCAG
 3101 CCACCCGCAA GGCTAACAAAC GACATTACTT CCATCACAAG CAACGGCGGA
 3151 AGAGTTAACT GCATGCAGGT GTGGCCTCCG ATTGGAAAGA AGAAGTTTGA
 3201 GACTCTCTCT TACCTTCCTG ACCTTACCGA TTCCGGTGGT CGCGTCAACT
 15 3251 GCATGCAGGC CATGGCTAAG ATTTTGTGATT TCGTAAAACC TGGCGTAATC
 3301 ACTGGTGTATG ACGTACAGAA AGTTTTCCAG GTAGCAAAAG AAAACAACCT
 3351 CGCACTGCCA GCAGTAAACT GCGTCGGTAC TGACTCCATC AACGCCGTAC
 3401 TGGAAACCGC TGCTAAAGTT AAAGCGCCGG TTATCGTTCA GTTCTCCAAC
 3451 GGTGGTGTCT CTTTATTCGC TGGTAAAGGC GTGAAATCTG ACGTTCCGCA
 20 3501 GGTGTGCTGT ATCCTGGGCG CGATCTCTGG TGCATCATC GTTACCAGA
 3551 TGGCTGAACA TTATGGTGT TCGGTATCC TGCACACTGA CCACTGCGCG
 3601 AAGAACTGC TGCCGTGGAT CGACGGTCTG TTGGACGCG GTGAAAAACA
 3651 CTTGCGAGCT ACCGGTAAGC CGCTGTTCTC TTCTCACATG ATCGACCTGT
 3701 CTGAAGAATC TCTGCAAGAG AACATCGAAA TCTGTCTTAA ATACCTGGAG
 25 3751 CGCATGTCCA AAATCGGCAT GACTCTGGAA ATCGAACTGG GTTGCACCGG
 3801 TGGTGAAGAA GACGGCGTGG ACAACAGCCA CATGGACGCT TCTGCACTGT
 3851 ACACCCAGCC GGAAGACGTT GATTACGCAT ACACCGAAT GAGCAAAATC
 3901 AGCCCGCGTT TCACCATCGC AGCGTCTTTC GGTAACGTAC ACGGTGTTTA
 3951 CAAGCCGGGT AACGTGGTTC TGACTCCGAC CATCCTGCGT GATTCTCAGG
 30 4001 AATATGTTTC CAAGAAACAC AACCTGCCGC ACAACAGCCT GAACTTCGTA
 4051 TTCCACGGTG GTTCCGGTTC TACTGCTCAG GAAATCAAAG ACTCCGTAAG
 4101 CTACGGCGTA GTAAAAATGA ACATCGATAC CGATACCCAA TGGGCAACCT
 4151 GGAAGGCGT TCTGAAGTAC TACAAAGCGA ACGAAGCTTA TCTGCAGGGT
 4201 CAGCTGGGTA ACCCGAAAGG CGAAGATCAG CCGAACAAGA AATACTACGA
 35 4251 TCCGCGCGTA TGGCTGCGTG CCGGTGAGAC TTCGATGATC GCTCGTCTGG
 4301 AGAAAGCATT CCAGGAACGT AACGCGATCG ACGTTCTGTA AGAGCTCGGT
 4351 ACCGGATCCA ATTCCCGATC GTTCAAACAT TTGGCAATAA AGTTTCTTAA
 4401 GATTGAATCC TGTGCGCGT CTTGCGATGA TTATCATATA ATTTCTGTTG
 4451 AATTACGTTA AGCATGTAAT AATTAACATG TAATGCATGA CGTTATTAT
 40 4501 GAGATGGGTT TTTATGATTA GAGTCCCGCA ATTATACATT TAATACGCGA
 4551 TAGAAAACAA AATATAGCGC GCAAAGTAGG ATAAATTATC GCGCGCGGTG
 4601 TCATCTATGT TACTAGATCG GGGATCGATC CCCGGGCGGC CGCCACTCGA
 4651 GTGGTGGCCG CATCGATCGT GAAGTTTCTC ATCTAAGCCC CCATTGCGAC
 4701 GTGAATGTAG ACACGTCGAA ATAAAGATTT CCGAATTAGA ATAATTTGTT
 45 4751 TATTGCTTTT GCCTATAAAT ACGACGGATC GTAATTTGTC GTTTTATCAA
 4801 AATGTAATTT CATTTTTATA TAACGCTGCG GACATCTACA TTTTGAATT
 4851 GAAAAAAAT TGGTAATTAC TCTTTCTTTT TCTCCATATT GACCATCATA
 4901 CTCATTGCTG ATCCATGTAG ATTTCCCGGA CATGAAGCCA TTTACAATTG
 4951 AATATATCCT GCCCGCGCTG CCGCTTTGCA CCCGGTGGAG CTTGCATGTT
 50 5001 GGTTCCTACG CAGAACTGAG CCGGTTAGGC AGATAATTC CATTGAGAAC
 5051 TGAGCCATGT GCACCTTCCC CCCAACACGG TGAGCGACGG GGCAACGGAG
 5101 TGATCCACAT GGGACTTTTC CTAGCTTGGC TGCCATTTT GGGGTGAGGC
 5151 CGTTCGCGCG GGGCGCCAGC TGGGGGGATG GGAGGCCCGC GTTACCGGGA
 5201 GGGTTCGAGA AGGGGGGGCA CCCCCCTTCG GCGTGCGCGG TCACGCGCCA
 55 5251 GGGCGCAGCC CTGGTTAAAA ACAAGGTTTA TAAATATTGG TTTAAAAAGCA
 5301 GGTTAAAAGA CAGGTTAGCG GTGGCCGAAA AACGGGCGGA AACCCCTGCA
 5351 AATGCTGGAT TTTCTGCCTG TGGACAGCCC CTCAAATGTC AATAGGTGCG
 5401 CCCCTCATCT GTCATCACTC TGCCCCCTCA GTGTCAAGGA TCGCGCCCCT

5451 CATCTGTCAG TAGTCGCGCC CCTCAAGTGT CAATACCGCA GGGCACTTAT
 5501 CCCAGGCTT GTCCACATCA TCTGTGGGAA ACTCGCGTAA AATCAGGCGT
 5551 TTTCGCGGAT TTGCGAGGCT GGCCAGCTCC ACGTGCGCCG CCGAAATCGA
 5601 GCCTGCCCCT CATCTGTCAA CGCCGCGCCG GGTGAGTCGG CCCCTCAAGT
 5 5651 GTCAACGTCC GCCCCTCATC TGTCAGTGAG GGCCAAGTTT TCCGCGTGGT
 5701 ATCCACAACG CCGGCGGCCG GCCGCGGTGT CTCGCACACG GCTTCGACGG
 5751 CGTTTCTGGC GCGTTTGACG GGCCATAGAC GGCCGCCAGC CCAGCGGCCA
 5801 GGGCAACCAG CCCGGTGAGC GTCGAAAGG GTCGATCGAC CGATGCCCTT
 5851 GAGAGCCTTC AACCAGTCA GCTCCTCCG GTGGGCGCGG GGCATGACTA
 10 5901 TCGTCGCCGC ACTTATGACT GTCTTCTTTA TCATGCAACT CGTAGGACAG
 5951 GTGCCGGCAG CGCTCTGGGT CATTTTCGGC GAGGACCGCT TTCGCTGGAG
 6001 CGCGACGATG ATCGGCCTGT CGTTGCGGT ATTCCGAATC TTGCACGCCC
 6051 TCGCTCAAGC CTTCTGCACT GGTCCCGCCA CCAAACGTTT CGGCGAGAAG
 6101 CAGGCCATTA TCGCCGGCAT GGCGGCCGAC GCGCTGGGCT ACGTCTTGCT
 15 6151 GCGGTTCCGCG ACGCGAGGCT GGATGGCCTT CCCCATATG ATTCTTCTCG
 6201 CTTCCGGCGG CATCGGGATG CCCGCGTTGC AGGCCATGCT GTCCAGGCAG
 6251 GTAGATGACG ACCATCAGGG ACAGCTTCAA GGATCGCTCG CGGCTCTTAC
 6301 CAGCCTAACT TCGATCACTG GACCGCTGAT CGTCACGGCG ATTTATGCCG
 6351 CCTCGGCGAG CACATGGAAC GGGTTGGCAT GGATTGTAGG CGCCGCCCTA
 20 6401 TACCTTGTCT GCCTCCCCGC GTTGCCTCGC GGTGCATGGA GCCGGGCCAC
 6451 CTCGACCTGA ATGGAAGCCG GCGGCACCTC GCTAACGGAT TCACCACTCC
 6501 AAGAATTGGA GCCAATCAAT TCTTGCGGAG AACTGTGAAT GCGCAAACCA
 6551 ACCCTTGGCA GAACATATCC ATCGCGTCCG CCATCTCCAG CAGCCGCACG
 6601 CGGCGCATCT CGGGCAGCGT TGGGTCCCTG CCACGGGTGC GCATGATCGT
 25 6651 GCTCCTGTCT TTGAGGACCC GGCTAGGCTG GCGGGGTTGC CTTACTGGTT
 6701 AGCAGAATGA ATCACCATA CGCGAGCGAA CGTGAAGCGA CTGCTGCTGC
 6751 AAAACGTCTG CGACCTGAGC AACAACATGA ATGGTCTTCG GTTCCGTGT
 6801 TTCGTAAAGT CTGGAACGC GGAAGTCAGC GCCCTGCACC ATTATGTTCC
 6851 GGATCTGCAT CGCAGGATGC TGCTGGCTAC CCTGTGGAAC ACCTACATCT
 30 6901 GTATTAACGA AGCGCTGGCA TTGACCCTGA GTGATTTTTC TCTGGTCCCG
 6951 CCGCATCCAT ACCGCCAGTT GTTTACCCTC ACAACGTTCC AGTAACCGGG
 7001 CATGTTTCATC ATCAGTAACC CGTATCGTGA GCATCCTCTC TCGTTTCATC
 7051 GGTATCATTA CCCCCATGAA CAGAAATTCC CCCTTACAG GAGGCATCAA
 7101 CCAGACAAAC AGGAAAAAAC CGCCCTTAA ATGGCCCGCT TTATCAGAAG
 35 7151 CCAGACATTA ACGCTTCTGG AGAAACTCAA CGAGCTGGAC GCGGATGAAC
 7201 AGGCAGACAT CTGTGAATCG CTTACGACC ACGCTGATGA GCTTTACCGC
 7251 AGCTGCCTCG CGCGTTTCGG TGATGACGGT GAAAACCTCT GACACATGCA
 7301 GCTCCCGGAG ACGGTCACAG CTTGTCTGTA AGCGGATGCC GGGAGCAGAC
 7351 AAGCCCGTCA GGGCGCGTCA GCGGGTGTG GCGGGTGTG GGGCGCAGCC
 40 7401 ATGACCCAGT CACGTAGCGA TAGCGGAGTG TATACTGGCT TAACTATGCG
 7451 GCATCAGAGC AGATTGTACT GAGAGTGAC CATATGCGGT GTGAAATACC
 7501 GCACAGATGC GTAAGGAGAA AATACCGCAT CAGGCGCTCT TCCGCTTCCT
 7551 CGCTCACTGA CTCGCTGCCG TCGGTCTGTC GGCTGCGCG AGCGGTATCA
 7601 GCTCACTCAA AGGCGGTAAT ACGGTTATCC ACAGAATCAG GGGATAACGC
 45 7651 AGGAAAGAAC ATGTGAGCAA AAGGCCAGCA AAAGGCCAGG AACCCTAAAA
 7701 AGGCCGCGTT GCTGGCGTTT TTCCATAGGC TCCGCCCCC TGACGAGCAT
 7751 CACAAAAATC GACGCTCAAG TCAGAGGTGG CGAAACCCGA CAGGACTATA
 7801 AAGATACCAG GCGTTTCCCC CTGGAAGCTC CCTCGTGCGC TCTCCTGTTT
 7851 CGACCTGCCC GCTTACCGGA TACCTGTCCG CCTTTCTCCC TTCGGGAAGC
 50 7901 GTGGCGCTTT CTCATAGCTC ACGCTGTAGG TATCTCAGTT CCGGTGTAGGT
 7951 CGTTCGCTCC AAGCTGGGCT GTGTGCACGA ACCCCCGTT CAGCCCGACC
 8001 GCTGCGCCTT ATCCGGTAAC TATCGTCTTG AGTCCAACCC GGTAAAGAC
 8051 GACTTATCGC CACTGGCAGC AGCCACTGGT AACAGGATTA GCAGAGCGAG
 8101 GTATGTAGGC GGTGCTACAG AGTTCTTGAA GTGGTGGCCT AACTACGGCT
 55 8151 AACTAGAAAG GACAGTATTT GGTATCTGCG CTCTGCTGAA GCCAGTTACC
 8201 TTCGGAAAAA GAGTTGGTAG CTCTTGATCC GGCAAAACAA CCACCGCTGG
 8251 TAGCGGTGGT TTTTGTGTTT GCAAGCAGCA GATTACGCGC AGAAAAAAG
 8301 GATCTCAAGA AGATCCTTTG ATCTTTTCTA CGGGGTCTGA CGCTCAGTGG

```

8351 AACGAAACT CACGTTAAGG GATTTTGGTC ATGAGATTAT CAAAAAGGAT
8401 CTTACCTTAG ATCCTTTTAA ATTAAAAATG AAGTTTAA TCAATCTAAA
8451 GTATATATGA GTAAACTTGG TCTGACAGTT ACCAATGCTT AATCAGTGAG
8501 GCACCTATCT CAGCGATCTG TCTATTTCTG TCATCCATAG TTGCCTGACT
5 8551 CCCCCTCGTG TAGATAACTA CGATACGGGA GGGCTTACCA TCTGGCCCCA
8601 GTGCTGCAAT GATACCGCGA GACCCACGCT CACCGGCTCC AGATTATCA
8651 GCAATAAACC AGCCAGCCGG AAGGGCCGAG CGCAGAAAGTG GTCCTGCAAC
8701 TTTATCCGCC TCCATCCAGT CTATTAATTG TTGCCGGGAA GCTAGAGTAA
8751 GTAGTTCGCC AGTTAATAGT TTGCGCAACG TTGTTGCCAT TGCTGCAGGT
10 8801 CGGGAGCACA GGATGACGCC TAACAATTCA TTCAAGCCGA CACCGCTTCG
8851 CGGCGCGGCT TAATTCAGGA GTTAAACATC ATGAGGGAAG CGGTGATCGC
8901 CGAAGTATCG ACTCAACTAT CAGAGGTAGT TGGCGTCATC GAGCGCCATC
8951 TCGAACCGAC GTTGCTGGCC GTACATTTGT ACGGCTCCGC AGTGGATGGC
9001 GGCCTGAAGC CACACAGTGA TATTGATTG CTGGTTACGG TGACCGTAAG
15 9051 GCTTGATGAA ACAACGCGC GAGCTTTGAT CAACGACCTT TTGGAAACTT
9101 CGGCTTCCCC TGGAGAGAGC GAGATTCTCC GCGCTGTAGA AGTCACCATT
9151 GTTGTGCACG ACGACATCAT TCCGTGGCGT TATCCAGCTA AGCGCGAACT
9201 GCAATTTGGA GAATGGCAGC GCAATGACAT TCTTGCAAGT ATCTTCGAGC
9251 CAGCCACGAT CGACATTGAT CTGGCTATCT TGCTGACAAA AGCAAGAGAA
20 9301 CATAGCGTTG CCTTGGTAGG TCCAGCGGCG GAGGAACTCT TTGATCCGGT
9351 TCCTGAACAG GATCTATTG AGGCGCTAAA TGAACCTTA ACGCTATGGA
9401 ACTCGCCGCC CGACTGGGCT GGCGATGAGC GAAATGTAGT GCTTACGTTG
9451 TCCCGCATTT GGTACAGCGC AGTAACCGGC AAAATCGCGC CGAAGGATGT
9501 CGCTGCCGAC TGGGCAATGG AGCGCCTGCC GGCCAGTAT CAGCCCCTCA
25 9551 TACTTGAAGC TAGGCAGGCT TATCTTGGAC AAGAAGATCG CTTGGCCTCG
9601 CGCGCAGATC AGTTGGAAGA ATTTGTTTAC TACGTGAAAG GCGAGATCAC
9651 CAAGGTAGTC GGCAAATAAT GTCTAACAAT TCGTTCAAGC CGACGCCGCT
9701 TCGCGGCGCG GCTTAACTCA AGCGTTAGAT GCTGCAGGCA TCGTGGTGTC
9751 ACGCTCGTCG TTTGGTATGG CTTCAATTCAG CTCCGGTTCC CAACGATCAA
30 9801 GGCAGAGTTAC ATGATCCCCC ATGTTGTGCA AAAAAGCGGT TAGCTCCTTC
9851 GGTCTCCGA TCGAGGATTT TTCGGCGCTG CGCTACGTCC GCKACCGCGT
9901 TGAGGGATCA AGCCACAGCA GCCCACCTCGA CCTCTAGCCG ACCCAGACGA
9951 GCCAAGGGAT CTTTTTGAA TGCTGCTCCG TCGTCAGGCT TTCCGACGTT
10001 TGGGTGGTTG AACAGAAGTC ATTATCGTAC GGAATGCCAA GCACTCCCCG
35 10051 GGGGAACCCCT GTGGTTGGCA TGCACATACA AATGGACGAA CGGATAAACC
10101 TTTTCACGCC CTTTAAATA TCCGTTATTC TAATAAACGC TCTTTTCTCT
10151 TAGGTTTACC CGCCAATATA TCCTGTCAA CACTGATAGT TTAACTGAA
10201 GGCGGGAAAC GACAATCTGA TCCCCATCAA GCTTGAGCTC AGGATTTAGC
10251 AGCATTCAG ATTGGTTCA ATCAACAAGG TACGAGCCAT ATCACTTTAT
40 10301 TCAAATTGGT ATCGCCAAAA CCAAGAAGGA ACTCCCATCC TCAAAGGTTT
10351 GTAAGGAAGA ATTCTCAGTC CAAAGCCTCA ACAAGGTCAG GGTACAGAGT
10401 CTCCAAACCA TTAGCCAAAA GCTACAGGAG ATCAATGAAG AATCTTCAAT
10451 CAAAGTAAAC TACTGTTCCA GCACATGCAT CATGGTCAGT AAGTTTCAGA
10501 AAAAGACATC CACCGAAGAC TTAAAGTTAG TGGGCATCTT TGAAAGTAAT
45 10551 CTTGTCAACA TCGAGCAGCT GGCTTGTGGG GACCAGACAA AAAAGGAATG
10601 GTGCAGAAAT GTTAGGCGCA CCTACCAAAA GCATCTTTGC CTTTATTGCA
10651 AAGATAAAGC AGATTCTCT AGTACAAGTG GGGAACAAAA TAACGTGGAA
10701 AAGAGCTGTC CTGACAGCCC ACTCACTAAT GCGTATGACG AACGCAGTGA
10751 CGACCACAAA AGAATCCCT CTATATAAGA AGGCATTCAT TCCCATTGA
50 10801 AGGATCATCA GATACTGAAC CAATCCTTCT AGAAGATCTA AGCTTAT

```

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- 55 (A) LENGTH: 10901 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

```

1   CGATAAGCTT GATGTAATTG GAGGAAGATC AAAATTTTCA ATCCCCATTC
5   51   TTCGATTGCT TCAATTGAAG TTTCTCCGAT GGCGCAAGTT AGCAGAATCT
    101  GCAATGGTGT GCAGAACCCA TCTCTTATCT CCAATCTCTC GAAATCCAGT
    151  CAACGCAAAT CTCCCTTATC GGTTCCTCTG AAGACGCAGC AGCATCCACG
    201  AGCTTATCCG ATTTTCGTCGT CGTGGGGATT GAAGAAGAGT GGGATGACGT
    251  TAATTGGCTC TGAGCTTCGT CCTCTTAAGG TCATGTCTTC TGTTCACAG
10   301  GCGTGCGATG TTCACGGTGC AAGCAGCCGT CCAGCAACTG CTCGTAAGTC
    351  CTCTGGTCTT TCTGGAACCG TCCGTATTCC AGGTGACAAG TCTATCTCCC
    401  ACAGGTCCTT CATGTTTGA GGTCTCGCTA GCGGTGAAAC TCGTATCACC
    451  GGTCTTTTGG AAGGTGAAGA TGTATCAAC ACTGGTAAGG CTATGCAAGC
    501  TATGGGTGCC AGAATCCGTA AGGAAGGTGA TACTTGATC ATTGATGGTG
15   551  TTGGTAACGG TGGACTCCTT GCTCCTGAGG CTCCTCTCGA TTTCGGTAAC
    601  GCTGCAACTG GTTGCCGTTT GACTATGGGT CTTGTTGGTG TTTACGATTT
    651  CGATAGCACT TTCATTGGTG ACGCTTCTCT CACTAAGCGT CCAATGGGTC
    701  GTGTGTTGAA CCCACTTCGC GAAATGGGTG TGCAGGTGAA GTCTGAAGAC
    751  GGTGATCGTC TTCCAGTTAC CTTGCGTGGA CCAAAGACTC CAACGCCAAT
20   801  CACCTACAGG GTACCTATGG CTTCCGCTCA AGTGAAGTCC GCTGTTCTGC
    851  TTGCTGGTCT CAACACCCCA GGTATCACCA CTGTTATCGA GCCAATCATG
    901  ACTCGTGACC AACTGAAAA GATGCTTCAA GGTTTTGGTG CTAACCTTAC
    951  CGTTGAGACT GATGCTGACG GTGTGCGTAC CATCCGTCTT GAAGGTCGTG
25   1001 GTAAGCTCAC CGGTCAAGTG ATTGATGTTT CAGGTGATCC ATCCTCTACT
    1051 GCTTTCCCAT TGGTTGCTGC CTTGCTTGTT CCAGGTTCCG ACGTCACCAT
    1101 CCTTAACGTT TTGATGAACC CAACCCGTAC TGGTCTCATC TTGACTCTGC
    1151 AGGAAATGGG TGCCGACATC GAAGTGATCA ACCCAGTCT TGCTGGTGGA
    1201 GAAGACGTGG CTGACTTGCG TGTTCGTTCT TCTACTTTGA AGGGTGTTAC
    1251 TGTTCCAGAA GACCGTGCTC CTTCTATGAT CGACGAGTAT CCAATTCTCG
30   1301 CTGTTGCAGC TGCATTGCTG GAAGGTGCTA CCGTTATGAA CGGTTTGAA
    1351 GAACTCCGTG TTAAGGAAAG CGACCGTCTT TCTGCTGTCG CAAACGGTCT
    1401 CAAGCTCAAC GGTGTTGATT GCGATGAAGG TGAGACTTCT CTCGTCGTGC
    1451 GTGGTCGTCC TGACGGTAAG GGTCTCGGTA ACGCTTCTGG AGCAGCTGTC
    1501 GCTACCCACC TCGATCACCG TATCGCTATG AGCTTCTCTG TTATGGGTCT
35   1551 CGTTTCTGAA AACCTGTTA CTGTTGATGA TGCTACTATG ATCGCTACTA
    1601 GCTTCCAGAA GTTCATGGAT TTGATGGCTG GTCTTGAGC TAAGATCGAA
    1651 CTCTCCGACA CTAAGGCTGC TTGATGAGCT CAAGAATTCT AGCTCGGTAC
    1701 CGGATCCAGC TTTTCGTTCT ATCATCGGTT TCGACAACGT TCGTCAAGTT
    1751 CAATGCATCA GTTTCATTGC GCACACACCA GAATCCTACT GAGTTCGAGT
40   1801 ATTATGGCAT TGGGAAAAC GTTTTTCTTG TACCATTTGT TGTGCTTGTA
    1851 ATTTACTGTG TTTTTTATTC GGTTTTCGCT ATCGAACTGT GAAATGGAAA
    1901 TGGATGGAGA AGAGTTAATG AATGATATGG TCCTTTTGTT CATTTCTCAA
    1951 TTAATATTAT TTGTTTTTTC TCTTATTGTT TGTGTGTTGA ATTTGAAATT
    2001 ATAAGAGATA TGCAAAACATT TTGTTTGTAG TAAAAATGTG TCAAATCGTG
45   2051 GCCTCTAATG ACCGAAGTTA ATATGAGGAG TAAAAACTTT GTAGTTGTAC
    2101 CATTATGCTT ATTCACTAGG CAACAAATAT ATTTTCAGAC CTAGAAAAGC
    2151 TGCAAATGTT ACTGAATACA AGTATGTCCT CTTGTGTTTT AGACATTTAT
    2201 GAACTTTTCT TTATGTAATT TTCCAGAATC CTTGTCAGAT TCTAATCATT
    2251 GCTTTATAAT TATAGTTATA CTCATGGATT TGTAGTTGAG TATGAAAATA
50   2301 TTTTTTAATG CATTTTATGA CTTGCCAATT GATTGACAAC ATGCATCAAT
    2351 CGACCTGCAG CCACTCGAAG CGGCCGCGTT CAAGCTTGAG CTCAGGATTT
    2401 AGCAGCATTC CAGATTGGGT TCAATCAACA AGGTACGAGC CATATCACTT
    2451 TATTCAAATT GGTATCGCCA AAACCAAGAA GGAACCTCCA TCCTCAAAGG
    2501 TTTGTAAGGA AGAATTCTCA GTCCAAAGCC TCAACAAGGT CAGGGTACAG
55   2551 AGTCTCCAAA CCATTAGCCA AAAGCTACAG GAGATCAATG AAGAATCTTC
    2601 AATCAAAGTA AACTACTGTT CCAGCACATG CATCATGGTC AGTAAGTTTC
    2651 AGAAAAAGAC ATCCACCGAA GACTTAAAGT TAGTGGGCAT CTTTGAAAGT
    2701 AATCTTGTC AATCGAGCA GCTGGCTTGT GGGGACCAGA CAAAAAAGGA

```

2751 ATGGTGCAGA ATTGTTAGGC GCACCTACCA AAAGCATCTT TGCCTTTATT
 2801 GCAAAGATAA AGCAGATTCC TCTAGTACAA GTGGGAACA AAATAACGTG
 2851 GAAAAGAGCT GTCCTGACAG CCCACTCACT AATGCGTATG ACGAACGCAG
 2901 TGACGACCAC AAAAGAATTC CCTCTATATA AGAAGGCATT CATTCCCATT
 5 2951 TGAAGGATCA TCAGATACTG AACCAATCCT TCTAGAAGAT CTAAGCTTAT
 3001 CGATAAGCTT GATGTAATTG GAGGAAGATC AAAATTTTCA ATCCCCATT
 3051 TTCGATTGCT TCAATTGAAG TTTCTCCGAT GGCACAAGTT AGCAGAATCT
 3101 GCAATGGTGT GCAGAACCCA TCTCTTATCT CCAATCTCTC GAAATCCAGT
 3151 CAACGCAAAT CTCCCTTATC GGTTCCTCTG AAGACGCAGC AGCATCCACG
 10 3201 AGCTTATCCG ATTTCTGTCG CTGCGGGATT GAAGAAGAGT GGGATGACGT
 3251 TAATTGGCTC TGAGCTTCGT CCTCTTAAGG TCATGTCTTC TGTTCACG
 3301 GCGTGCATGC AGGCcatggC TAAGATTTTT GATTCGTAA AACCTGGCGT
 3351 AATCACTGGT GATGACGTAC AGAAAGTTTT CCAGGTAGCA AAAGAAAACA
 3401 ACTTCGCACT GCCAGCAGTA AACTGCGTCG GTACTGACTC CATCAACGCC
 15 3451 GTACTGGAAA CCGCTGCTAA AGTTAAAGCG CCGGTATCG TTCAGTTCTC
 3501 CAACGGTGGT GCTTCCTTTA TCGCTGGTAA AGGCGTGAAA TCTGACGTTT
 3551 CGCAGGGTGC TGCTATCCTG GCGCGGATCT CTGGTGCGCA TCACGTTTAC
 3601 CAGATGGCTG AACATTATGG TGTTCGGTT ATCCTGCACA CTGACCACTG
 3651 CCGGAAGAAA CTGCTGCCGT GGATCGACGG TCTGTTGGAC GCGGGTGAAA
 20 3701 AACACTTCGC AGCTACCGGT AAGCCGCTGT TCTCTTCTCA CATGATCGAC
 3751 CTGTCTGAAG AATCTCTGCA AGAGAACATC GAAATCTGCT CTAAATACCT
 3801 GGAGCGCATG TCCAAAATCG GCATGACTCT GGAAATCGAA CTGGGTTGCA
 3851 CCGGTGGTGA AGAAGACGGC GTGGACAACA GCCACATGGA CGCTTCTGCA
 3901 CTGTACACCC AGCCGGAAGA CGTTGATTAC GCATACACCG AACTGAGCAA
 25 3951 AATCAGCCCG CGTTTCACCA TCGCAGCGTC CTTCCGTAAC GTACACGGTG
 4001 TTTACAAGCC GGGTAACGTG GTTCTGACTC CGACCATCCT GCGTGATTCT
 4051 CAGGAATATG TTTCCAAGAA ACACAACCTG CCGACAACA GCCTGAACCT
 4101 CGTATTCCAC GGTGGTTCCG GTTCTACTGC TCAGGAAATC AAAGACTCCG
 4151 TAAGCTACGG CGTAGTAAAA ATGAACATCG ATACCGATAC CCAATGGGCA
 30 4201 ACCTGGGAAG GCGTCTGAA CTACTACAAA GCGAACGAAG CTTATCTGCA
 4251 GGGTCAGCTG GGTAACCCGA AAGGCGAAGA TCAGCCGAAC AAGAAATACT
 4301 ACGATCCCGC CGTATGGCTG CGTGCCGGTC AGACTTCGAT GATCGCTCGT
 4351 CTGGAGAAAG CATTCCAGGA ACTGAACGCG ATCGACGTTT TGTAAAGACT
 4401 CCGTACCGGA TCCAATTccc GATCGTTCAA ACATTGGCA ATAAAGTTTC
 35 4451 TTAAGATTGA ATCCTGTTGC CGGTCTTGC ATGATTATCA TATAATTCT
 4501 GTTGAATTAC GTTAAGCATG TAATAATTAA CATGTAATGC ATGACGTTAT
 4551 TTATGAGATG GGTTCCTATG ATTAGAGTCC CGCAATTATA CATTTAATAC
 4601 GCGATAGAAA ACAAATATA GCGCGCAAAC TAGGATAAAT TATCGCGCGC
 4651 GGTGTCATCT ATGTTACTAG ATCGGGGATC GATCCCCGGG CCGCCGCCAC
 40 4701 TCGAGTGGTG GCCGCATCGA TCGTGAAGTT TCTCATCTAA GCCCCATT
 4751 GGACGTGAAT GTAGACACGT CGAAATAAAG ATTTCCGAAT TAGAATAATT
 4801 TGTTTATTGC TTTCGCCCTAT AAATACGACG GATCGTAATT TGTCGTTTTA
 4851 TCAAAATGTA CTTTCATTTT ATAATAACGC TCGGACATC TACATTTTG
 4901 AATTGAAAAA AAATTGGTAA TTAATCTTTC TTTTCTCCA TATTGACCAT
 45 4951 CATACTCATT GCTGATCCAT GTAGATTTC CGGACATGAA GCCATTTACA
 5001 ATTGAATATA TCCTGCCGCC GCTGCCGCTT TGCACCCGGT GGAGCTTGCA
 5051 TGTTGGTTTC TACGCAGAAC TGAGCCGCTT AGGCAGATAA TTCCATTGA
 5101 GAACTGAGCC ATGTGCACCT TCCCCCAAC ACGGTGAGCG ACGGGGCAAC
 5151 GGAAGTATCC ACATGGGACT TTTCCTAGCT TGGCTGCCAT TTTTGGCGTG
 50 5201 AGGCCGTTTC GCGGGGGCGC CAGCTGGGGG GATGGGAGGC CCGCGTTACC
 5251 GGGAGGGTTC GAGAAGGGGG GGCACCCCCC TTCGCGTGC GCGGTACGC
 5301 GCCAGGGCGC AGCCCTGGTT AAAAAACAAGG TTTATAAATA TTGGTTTAAA
 5351 AGCAGGTTAA AAGACAGGTT AGCGGTGGCC GAAAAACGGG CGGAAACCCT
 5401 TGCAAATGCT GGATTTTCTG CCTGTGGACA GCCCCTCAA TGTCAATAGG
 55 5451 TGCGCCCTC ATCTGTATC ACTCTGCCCC TCAAGTGTCA AGGATCGCGC
 5501 CCCTCATCTG TCAGTAGTCG CGCCCTCAA GTGTCAATAC CGCAGGGCAC
 5551 TTATCCCCAG GCTTGTCCAC ATCATCTGTG GGAACTCGC GTAAATCAG
 5601 GCGTTTTTCG CGATTTGCGA GGCTGGCCAG CTCCACGTG CCGCCGAA

	5651	TCGAGCCTGC	CCCTCATCTG	TCAACGCCGC	GCCGGGTGAG	TCGGCCCCTC
	5701	AAGTGTCAAC	GTCCGCCCCCT	CATCTGTCTAG	TGAGGGCCAA	GTTTTCCGCG
	5751	TGGTATCCAC	AACGCCGGCG	GCCGGCCGCG	GTGTCTCGCA	CACGGCTTCG
	5801	ACGGCGTTTT	TGGCGCGTTT	GCAGGGCCAT	AGACGGCCGC	CAGCCCAGCG
5	5851	GCGAGGGCAA	CCAGCCCCGT	GAGCGTCGGA	AAGGGTCGAT	CGACCGATGC
	5901	CCTTGAGAGC	CTTCAACCCA	GTCAGCTCCT	TCCGGTGGGC	GCGGGGCATG
	5951	ACTATCGTCG	CCGCACTTAT	GACTGTCTTC	TTTATCATGC	AACTCGTAGG
	6001	ACAGGTGCCG	GCAGCGCTCT	GGGTCATTTT	CGGCGAGGAC	CGCTTTCGCT
	6051	GGAGCGCGAC	GATGATCGGC	CTGTCTGCTT	CGGTATTCGG	AATCTTGCAC
10	6101	GCCCTCGCTC	AAGCCTTCGT	CACTGGTCCC	GCCACCAAAC	GTTTCGGCGA
	6151	GAAGCAGGCC	ATTATCGCCG	GCATGGCGGC	CGACGCGCTG	GGCTACGTCT
	6201	TGCTGGCGTT	CGCGACGCGA	GGCTGGATGG	CCTTCCCCAT	TATGATTCTT
	6251	CTCGCTTCCG	GCGGCATCGG	GATGCCCGCG	TTGCAGGCCA	TGCTGTCCAG
	6301	GCAGGTAGAT	GACGACCATC	AGGGACAGCT	TCAAGGATCG	CTCGCGGCTC
15	6351	TTACCAGCCT	AACTTCGATC	ACTGGACCGC	TGATCGTCAC	GGCGATTTAT
	6401	GCCGCTTCGG	CGAGCACATG	GAACGGGTG	GCATGGATTG	TAGGCGCCGC
	6451	CCTATACCTT	GTCTGCCTCC	CCGCGTTGCG	TCGCGGTGCA	TGGAGCCGGG
	6501	CCACCTCGAC	CTGAATGGAA	GCCGGCGGCA	CCTCGCTAAC	GGATTACCA
	6551	CTCCAAGAAT	TGGAGCCAAT	CAATTCTTGC	GGAGAAGTGT	GAATGCGCAA
20	6601	ACCAACCCTT	GGCAGAACAT	ATCCATCGCG	TCCGCCATCT	CCAGCAGCCG
	6651	CACGCGGCGC	ATCTCGGGCA	GCGTTGGGTC	CTGGCCACGG	GTGCGCATGA
	6701	TCGTGCTCCT	GTCGTTGAGG	ACCCGGCTAG	GCTGGCGGGG	TTGCCTTACT
	6751	GGTTAGCAGA	ATGAATCACC	GATACGCGAG	CGAACGTGAA	GCGACTGCTG
	6801	CTGCAAAACG	TCTGCGACCT	GAGCAACAAC	ATGAATGGTC	TTCGGTTTCC
25	6851	GTGTTTCGTA	AAGTCTGGAA	ACGCGGAAGT	CAGCGCCCTG	CACCATTATG
	6901	TTCCGGATCT	GCATCGCAGG	ATGCTGCTGG	CTACCCTGTG	GAACACCTAC
	6951	ATCTGTATTA	ACGAAGCGCT	GGCATTGACC	CTGAGTGATT	TTTCTCTGCT
	7001	CCCGCCGCAT	CCATACCGCC	AGTTGTTTAC	CCTCACAACG	TTCCAGTAAC
	7051	CGGGCATGTT	CATCATCAGT	AACCCGTATC	GTGAGCATCC	TCTCTCGTTT
30	7101	CATCGGTATC	ATTACCCCCA	TGAACAGAAA	TTCCCCCTTA	CACGGAGGCA
	7151	TCAAGTGACC	AAACAGGAAA	AAACCGCCCT	TAACATGGCC	CGCTTTATCA
	7201	GAAGCCAGAC	ATTAACGCTT	CTGGAGAAAC	TCAACGAGCT	GGACGCGGAT
	7251	GAACAGGCAG	ACATCTGTGA	ATCGCTTCAC	GACCACGCTG	ATGAGCTTTA
35	7301	CCGCACTGTC	CTCGCGCGTT	TCGGTGATGA	CGGTGAAAAC	CTCTGACACA
	7351	TGCAGTCCCC	GGAGACGGTC	ACAGCTTGTC	TGTAAGCGGA	TGCCGGGAGC
	7401	AGACAAGCCC	GTCAGGGCGC	GTCAGCGGGT	GTTGGCGGGT	GTCGGGGCGC
	7451	AGCCATGACC	CAGTCACGTA	GCGATAGCGG	AGTGTATACT	GGCTTAACTA
	7501	TGCGGCATCA	GAGCAGATTG	TACTGAGAGT	GCACCATATG	CGGTGTGAAA
	7551	TACCGCACAG	ATGCGTAAGG	AGAAAATACC	GCATCAGGCG	CTCTTCCGCT
40	7601	TCCTCGCTCA	CTGACTCGCT	GCGCTCGGTC	GTTGCGCTGC	GGCGAGCGGT
	7651	ATCAGCTCAC	TCAAAGGCGG	TAATACGGTT	ATCCACAGAA	TCAGGGGATA
	7701	ACGCAGGAAA	GAACATGTGA	GCAAAAGGCC	AGCAAAAGGC	CAGGAACCGT
	7751	AAAAAGGCCG	CGTTGCTGGC	GTTTTTCCAT	AGGCTCCGCC	CCCCTGACGA
	7801	GCATCACAAA	AATCGACGCT	CAAGTCAGAG	GTGGCGAAAC	CCGACAGGAC
45	7851	TATAAAGATA	CCAGGCGTTT	CCCCCTGGAA	GCTCCCTCGT	GCGCTCTCCT
	7901	GTTCCGACCC	TGCCGCTTAC	CGGATACCTG	TCCGCCTTTG	TCCCTTCGGG
	7951	AAGCGTGGCG	CTTCTCTATA	GCTCACGCTG	TAGGTATCTC	AGTTCGGTGT
	8001	AGGTCGTTTC	CTCCAAGCTG	GGCTGTGTGC	ACGAACCCCC	CGTTCAGCCC
	8051	GACCGCTGCG	CCTTATCCGG	TAACTATCGT	CTTGAGTCCA	ACCCGGTAAG
50	8101	ACACGACTTA	TCGCCACTGG	CAGCAGCCAC	TGGTAACAGG	ATTAGCAGAG
	8151	CGAGGTATGT	AGGCGGTGCT	ACAGAGTTCT	TGAAGTGGTG	GCCTAACTAC
	8201	GGCTACACTA	GAAGGACAGT	ATTTGGTATC	TGCGCTCTGC	TGAAGCCAGT
	8251	TACCTTCGGA	AAAAGAGTTG	GTAGCTCTTG	ATCCGGCAAA	CAAAACCACG
	8301	CTGGTAGCGG	TGGTTTTTTT	GTTTGCAAGC	AGCAGATTAC	GCGCAGAAAA
55	8351	AAAGGATCTC	AAGAAGATCC	TTTGATCTTT	TCTACGGGGT	CTGACGCTCA
	8401	GTGGAACGAA	AACTCACGTT	AAGGGATTTT	GGTCATGAGA	TTATCAAAAA
	8451	GGATCTTCAC	CTAGATCCTT	TTAAATTAAA	AATGAAGTTT	TAAATCAATC
	8501	TAAAGTATAT	ATGAGTAAAC	TTGGTCTGAC	AGTTACCAAT	GCTTAATCAG

8551 TGAGGCACCT ATCTCAGCGA TCTGTCTATT TCGTTCATCC ATAGTTGCCT
 8601 GACTCCCCGT CGTG TAGATA ACTACGATAC GGGAGGGCTT ACCATCTGGC
 8651 CCCAGTGCTG CAATGATACC GCGAGACCCA CGCTCACC GG CTCCAGATTT
 8701 ATCAGCAATA AACCAGCCAG CCGGAAGGGC CGAGCGCAGA AGTGGTCTTG
 5 8751 CAACTTTATC CGCCTCCATC CAGTCTATTA ATTGTTGCCG GGAAGCTAGA
 8801 GTAAGTAGTT CGCCAGTTAA TAGTTTGCGC AACGTTGTTG CCATTGCTGC
 8851 AGGTCGGGAG CACAGGATGA CGCCTAACAA TTCATTCAAG CCGACACCGC
 8901 TTCGCGGCGC GGCTTAATTC AGGAGTTAAA CATCATGAGG GAAGCGGTGA
 8951 TCGCCGAAGT ATCGACTCAA CTATCAGAGG TAGTTGGCGT CATCGAGCGC
 10 9001 CATCTCGAAC CGACGTTGCT GGCCGTACAT TTGTACGGCT CCGCAGTGGA
 9051 TGGCGGCCCTG AAGCCACACA GTGATATTGA TTTGCTGGTT ACGGTGACCG
 9101 TAAGGCTTGA TGAAACAACG CGGCGAGCTT TGATCAACGA CCTTTTGGA
 9151 ACTTCGGCTT CCCCTGGAGA GAGCGAGATT CTCGCGCTG TAGAAGTCAC
 9201 CATTGTTGTG CACGACGACA TCATTCCGTG GCGTTATCCA GCTAAGCGCG
 15 9251 AACTGCAATT TGGAGAATGG CAGCGCAATG ACATTCTTGC AGGTATCTTC
 9301 GAGCCAGCCA CGATCGACAT TGATCTGGCT ATCTTGCTGA CAAAAGCAAG
 9351 AGAACATAGC GTTGCCCTTG TAGGTCCAGC GCGGAGGAA CTCTTTGATC
 9401 CGGTTCTCTGA ACAGGATCTA TTTGAGGCGC TAAATGAAAC CTTAACGCTA
 9451 TGGAACTCGC CGCCCGACTG GGCTGGCGAT GAGCGAAATG TAGTGCTTAC
 20 9501 GTTGTCCTCGC ATTTGGTACA GCGCAGTAAC CGGCAAAATC GCGCCGAAGG
 9551 ATGTCGCTGC CGACTGGGCA ATGGAGCGCC TGCCGGCCCA GTATCAGCCC
 9601 GTCATACTTG AAGCTAGGCA GGCTTATCTT GGACAAGAAG ATCGCTTGCG
 9651 CTCGCGCGCA GATCAGTTGG AAGAATTTGT TCACTACGTG AAAGGCGAGA
 9701 TCACCAAGGT AGTCGGCAAA TAATGTCTAA CAATTCGTTT AAGCCGACGC
 25 9751 CGCTTCGCGG CGCGGCTTAA CTCAAGCGTT AGATGCTGCA GGCATCGTGG
 9801 TGTCACGCTC GTCGTTTGGT ATGGCTTCAT TCAGCTCCGG TTCCCAACGA
 9851 TCAAGGCGAG TTACATGATC CCCCATGTTG TGCAAAAAAG CGGTTAGCTC
 9901 CTTGCTGCTC CCGATCGAGG ATTTTTCGGC GCTGCGCTAC GTCCGCKACC
 9951 GCTTGAGGG ATCAAGCCAC AGCAGCCAC TCGACCTCTA GCCGACCCAG
 30 10001 ACGAGCCAAG GGATCTTTTT GGAATGCTGC TCCGTCGTCA GGCTTTCCGA
 10051 CGTTTGGGTG GTTGAACAGA AGTCATTATC GTACGGAATG CCAAGCACTC
 10101 CCGAGGGGAA CCCTGTGGTT GGCATGCACA TACAAATGGA CGAACGGATA
 10151 AACCTTTTCA CGCCCTTTTA AATATCCGTT ATTCTAATAA ACGCTCTTTT
 10201 CTCTTAGGTT TACCCGCCAA TATATCCTGT CAAACACTGA TAGTTTAAAC
 35 10251 TGAAGGCGGG AAACGACAAT CTGATCCCCA TCAAGCTTGA GCTCAGGATT
 10301 TAGCAGCATT CCAGATTGGG TTCAATCAAC AAGGTACGAG CCATATCACT
 10351 TTATTCAAAT TGGTATCGCC AAAACCAAGA AGGAACTCCC ATCCTCAAAG
 10401 GTTTGTAAAG AAGAATTCTC AGTCCAAAGC CTCAACAAGG TCAGGGTACA
 10451 GAGTCTCCAA ACCATTAGCC AAAAGCTACA GGAGATCAAT GAAGAATCTT
 40 10501 CAATCAAAGT AACTACTGT TCCAGCACAT GCATCATGGT CAGTAAGTTT
 10551 CAGAAAAAGA CATCCACCGA AGACTTAAAG TTAGTGGGCA TCTTTGAAAG
 10601 TAATCTTGTC AACATCGAGC AGCTGGCTTG TGGGGACCAG ACAAAAAAGG
 10651 AATGGTGCAG AATTGTTAGG CGCACCTACC AAAAGCATCT TTGCCTTTAT
 10701 TGCAAGATA AAGCAGATTC CTCTAGTACA AGTGGGGAAC AAAATAACGT
 45 10751 GGAAAAGAGC TGTCTGACA GCCCACTCAC TAATGCGTAT GACGAACGCA
 10801 GTGACGACCA CAAAAGAATT CCTCTATAT AAGAAGGCAT TCATTCCCAT
 10851 TTGAAGGATC ATCAGATACT GAACCAATCC TTCTAGAAGA TCTAAGCTTA
 10901 T

50

CLAIMS

1. A recombinant, double-stranded DNA molecule containing
 - a) a promoter functional in plant cells, and
 - 5 b) a DNA sequence coding for a polypeptide having the enzymatic activity of a fructose-1,6-bisphosphate aldolase and operatively linked to the promoter in sense orientation.
2. The DNA molecule according to claim 1, wherein the DNA sequence coding for a polypeptide having the enzymatic activity of a fructose-1,6-
10 bisphosphate aldolase is derived from a prokaryotic organism.
3. The DNA molecule according to claim 2, wherein the prokaryotic organism is *Escherichia coli*.
4. The DNA molecule according to claim 1, wherein the DNA sequence coding for a polypeptide having the enzymatic activity of a fructose-1,6-bisphosphate aldolase
15 has at least about 60% identity with a prokaryotic DNA sequence coding for fructose-1,6-bisphosphate aldolase class II.
5. The DNA molecule according to claim 1, wherein the DNA sequence coding for the polypeptide having the enzymatic activity of a fructose-1,6-bisphosphate aldolase is a sequence capable of hybridizing with the coding region depicted as
20 SEQ ID NO. 1.
6. The DNA molecule according to claim 1, wherein the DNA sequence coding for a polypeptide having the enzymatic activity of a fructose-1,6-bisphosphate aldolase has at least about 60% identity with the coding region depicted as SEQ ID NO. 1.
7. The DNA molecule according to claim 1, wherein the DNA sequence coding for a
25 polypeptide having the enzymatic activity of a fructose-1,6-bisphosphate aldolase has at least about 70% identity with the coding region depicted as SEQ ID NO. 1.

8. The DNA molecule according to claim 1, wherein the DNA sequence coding for a polypeptide having the enzymatic activity of a fructose-1,6-bisphosphate aldolase has at least about 80% identity with the coding region depicted as SEQ ID NO. 1.
9. The DNA molecule according to claim 1, wherein the DNA sequence coding for the polypeptide having the enzymatic activity of a fructose-1,6-bisphosphate aldolase has the coding region depicted as SEQ ID NO. 1, or encodes the same peptide as SEQ ID NO. 1 in accordance with the degeneracy of the genetic code.
10. A transgenic plant cell containing in its genome a recombinant DNA molecule according to any of claims 1-9.
11. A transgenic plant containing plant cells according to claim 10.
12. The transgenic plant of claim 11, wherein the plant exhibits a property selected from the group consisting of increased photosynthesis rates, increased yields, increased growth rates and improved solids uniformity compared with plants that do not contain the recombinant DNA molecule.
13. The transgenic plant according to claim 11, which is a crop plant.
14. The transgenic plant according to claim 11, selected from the group consisting of corn, wheat, rice, tomato, potato, carrots, sweet potato, yams, artichoke, alfalfa, peanut, barley, cotton, soybean, canola, sunflower, sugarbeet, apple, pear, orange, peach, sugarcane, strawberry, raspberry, banana, grape, plantain, tobacco, lettuce, cassava, cruciferous vegetables, forestry species and horticultural species.
15. The transgenic plant of claim 11, wherein the plant is a potato.
16. A food product derived from the potato of claim 15.
17. The food product of claim 16, which is a french fry or a potato chip.

18. Propagation material derived from the transgenic plant of claim 11.
19. A process for increasing the photosynthesis rate in plants which comprises transforming plant cells with a DNA molecule according to any one of claims 1 to 9, and regenerating the transformed cells to produce a transgenic plant.
- 5 20. A process for increasing the yield in plants which comprises transforming plant cells with a DNA molecule according to any one of claims 1 to 9, and regenerating the transformed cells to produce a transgenic plant.
21. A process for increasing the growth rate in plants which comprises transforming plant cells with a DNA molecule according to any one of claims 1 to 9, and
10 regenerating the transformed cells to produce a transgenic plant.
22. A process for improving the solids uniformity in plants which comprises transforming plant cells with a DNA molecule according to any one of claims 1 to 9, and regenerating the transformed cells to produce a transgenic plant.
23. In a method for the processing of potatoes into fries or chips, the improvement
15 comprising, utilizing a potato that overexpresses the *fda* transgene providing a higher solids uniformity in such potato.

1/11

TGCAACTTGAAGTATGACGAGTATAAGGCCCGACGATACAGGACAAGAGACATGTCT
AAG MetSerLys

ATTTTGGATTTCGTAAAACCTGGCGTAATCACTGGTGATGACGTACAGAAAGTTTCCAG
IlePheAspPheValLysProGlyValIleThrGlyAspAspValGlnLysValPheGln

GTAGCAAAAGAAAACAACCTTCGCACTGCCAGCAGTAACTGCGTCGGTACTGACTCCATC
ValAlaLysGluAsnAsnPheAlaLeuProAlaValAsnCysValGlyThrAspSerIle

AACGCCGTACTGGAAACCGCTGCTAAAGTTAAAGCGCCGGTTATCGTTCAGTTCTCCAAC
AsnAlaValLeuGluThrAlaAlaLysValLysAlaProValIleValGlnPheSerAsn

GGTGGTGCTTCCTTTATCGCTGGTAAAGGCGTGAAATCTGACGTTCCGCAGGGTGCTGCT
GlyGlyAlaSerPheIleAlaGlyLysGlyValLysSerAspValProGlnGlyAlaAla

ATCCTGGGCGCGATCTCTGGTGCGCATCACGTTACCAGATGGCTGAACATTATGGTGTT
IleLeuGlyAlaIleSerGlyAlaHisHisValHisGlnMetAlaGluHisTyrGlyVal

CCGGTTATCCTGCACACTGACCACTGCGCGAAGAACTGCTGCCGTGGATCGACGGTCTG
ProValIleLeuHisThrAspHisCysAlaLysLysLeuLeuProTrpIleAspGlyLeu

TTGGACGCGGGTGAAAAACACTTCGCAGCTACCGGTAAGCCGCTGTTCTTCTCACATG
LeuAspAlaGlyGluLysHisPheAlaAlaThrGlyLysProLeuPheSerSerHisMet

ATCGACCTGTCTGAAGAATCTCTGCAAGAGAACATCGAAATCTGCTCTAAATACCTGGAG
IleAspLeuSerGluGluSerLeuGlnGluAsnIleGluIleCysSerLysTyrLeuGlu

CGCATGTCCAAAATCGGCATGACTCTGGAAATCGAACTGGGTTGCACCGGTGGTGAAGAA
ArgMetSerLysIleGlyMetThrLeuGluIleGluLeuGlyCysThrGlyGlyGluGlu

GACGGCGTGGACAACAGCCACATGGACGCTTCTGCACTGTACACCCAGCCGGAAGACGTT
AspGlyValAspAsnSerHisMetAspAlaSerAlaLeuTyrThrGlnProGluAspVal

GATTACGCATACACCGAACTGAGCAAAATCAGCCCGCGTTTCACCATCGCAGCGTCCTTC
AspTyrAlaTyrThrGluLeuSerLysIleSerProArgPheThrIleAlaAlaSerPhe

FIG. 1A

2/11

GGTAACGTACACGGTGTTTACAAGCCGGTAACGTGGTTCTGACTCCGACCATCCTGCGT
GlyAsnValHisGlyValTyrLysProGlyAsnValValLeuThrProThrIleLeuArg

GATTCTCAGGAATATGTTTCCAAGAAACACAACCTGCCGCACAACAGCCTGAACTTCGTA
AspSerGlnGluTyrValSerLysLysHisAsnLeuProHisAsnSerLeuAsnPheVal

TTCCACGGTGGTTCCGGTTCTACTGCTCAGGAAATCAAAGACTCCGTAAGCTACGGCGTA
PheHisGlyGlySerGlySerThrAlaGlnGluIleLysAspSerValSerTyrGlyVal

GTAAAAATGAACATCGATACCGATACCCAATGGGCAACCTGGGAAGGCGTTCTGAACTAC
ValLysMetAsnIleAspThrAspThrGlnTrpAlaThrTrpGluGlyValLeuAsnTyr

TACAAAGCGAACGAAGCTTATCTGCAGGGTCAGCTGGGTAACCCGAAAGGCGAAGATCAG
TyrLysAlaAsnGluAlaTyrLeuGlnGlyGlnLeuGlyAsnProLysGlyGluAspGln

CCGAACAAGAAATACTACGATCCGCGCGTATGGCTGCGTGCCGGTCAGACTTCGATGATC
ProAsnLysLysTyrTyrAspProArgValTrpLeuArgAlaGlyGlnThrSerMetIle

CCT GCTCGTCTGGAGAAAGCATTCCAGGAACTGAACGCGATCGACGTTCTGTAAGATATT
AlaArgLeuGluLysAlaPheGlnGluLeuAsnAlaIleAspValLeuEnd
TTCTGCTTATCTCAAGGCCCGCTCTGCGGGTCTTTTTTTCG

FIG. 1B

3/11




FIG. 2

4/11




FIG. 3

5/11




FIG. 4A




FIG. 4B

6/11




FIG. 4C

7/11




FIG. 5

8/11




FIG. 6

9/11




FIG. 7

10/11




FIG. 8

11/11




FIG. 9

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/12447

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C12N15/82 C12N15/60 A01H5/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N A01H

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	SONNEWALD, U., ET AL.: "Manipulation of sink-source relations in transgenic plants" PLANT CELL AND ENVIRONMENT, vol. 17, 1994, pages 649-658, XP002080527 cited in the application see page 651, right-hand column ---	1-23
A	WO 96 21737 A (INST GENBIOLOGISCHE FORSCHUNG ;SONNEWALD UWE (DE); KOSSMANN JENS ()) 18 July 1996 see the whole document ---	1-23
A	WO 96 24679 A (MONSANTO CO) 15 August 1996 see page 20, line 15 - page 22, line 17 ---	1-23
-/--		



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

13 October 1998

Date of mailing of the international search report

23/10/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.
Fax: (+31-70) 340-3016

Authorized officer

Maddox, A

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/12447

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	KOSSMANN J ET AL: "REDUCTION OF THE CHLOROPLASTIC FRUCTOSE-1,6-BISPHOSPHATASE IN TRANSGENIC POTATO PLANTS IMPAIRS PHOTOSYNTHESIS AND PLANT GROWTH" PLANT JOURNAL, vol. 6, no. 5, 1 November 1994, pages 637-650, XP002001924 see the whole document ---	1-23
A	ZRENNER, R., ET AL.: "Reduction of the cytosolic fructose-1,6-bisphosphatase in transgenic potato plants limits photosynthetic sucrose biosynthesis with no impact on plant growth and tuber yield" THE PLANT JOURNAL, vol. 9, no. 5, 1996, pages 671-681, XP002080528 see the whole document ---	1-23
A	SCOTT, P., ET AL.: "Carbon metabolism in leaves of transgenic tobacco (Nicotiana tabacum L.) containing elevated fructose 2,6-bisphosphate levels" THE PLANT JOURNAL, vol. 7, no. 3, 1995, pages 461-469, XP002080529 see the whole document ---	1-23
A	SCOTT, P., ET AL.: "Influence of elevated fructose-2,6-bisphosphate levels on starch metabolism in transgenic tobacco leaves in the dark" PLANT PHYSIOLOGY, vol. 108, 1995, XP002080530 see the whole document ---	1-23
A	JUAN A S ET AL: "OVEREXPRESSION OF CYTOSOLIC FRUCTOSE- 1,6 BISPHOSPHATASE IN TRANSGENIC TOBACCO PLANTS" PLANT PHYSIOLOGY, vol. 105, no. 1, 1 May 1994, page 118 XP002001923 see the whole document ---	1-23
A	WO 91 19806 A (MONSANTO CO) 26 December 1991 see the whole document -----	1-23

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 98/12447

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9621737 A	18-07-1996	DE 19502053 A	18-07-1996
		AU 4485896 A	31-07-1996
		CA 2209932 A	18-07-1996
		EP 0802982 A	29-10-1997
		HU 9800420 A	29-06-1998
WO 9624679 A	15-08-1996	AU 695131 B	06-08-1998
		AU 4923596 A	27-08-1996
		CA 2212560 A	15-08-1996
		CZ 9702543 A	14-01-1998
		EP 0808371 A	26-11-1997
		PL 321738 A	22-12-1997
		SK 108997 A	06-05-1998
		US 5716837 A	10-02-1998
WO 9119806 A	26-12-1991	AU 644203 B	02-12-1993
		AU 8220291 A	07-01-1992
		EP 0536293 A	14-04-1993
		FI 925228 A	18-11-1992
		US 5498830 A	12-03-1996
		US 5608149 A	04-03-1997

THIS PAGE BLANK (USPTO)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)